

The Riemann zeros as spectrum and the Riemann hypothesis

Germán Sierra

Instituto de Física Teórica UAM/CSIC,
Universidad Autónoma de Madrid, Cantoblanco, Madrid, Spain.

We present a spectral realization of the Riemann zeros based on the propagation of a massless Dirac fermion in a region of Rindler spacetime and under the action of delta function potentials localized on the square free integers. The corresponding Hamiltonian admits a self-adjoint extension that is tuned to the phase of the zeta function, on the critical line, in order to obtain the Riemann zeros as bound states. The model provides a proof of the Riemann hypothesis in the limit where the potentials vanish. Finally, we propose an interferometer that may yield an experimental observation of the Riemann zeros.

Contents

I. Introduction	2
II. The semiclassical XP Berry, Keating and Connes model	2
III. The quantum XP model	4
IV. The Landau model and XP	5
V. The XP model revisited	7
VI. The space-time geometry of the modified XP models	10
VII. Diracization of $H = X(P + \ell_p^2/P)$	12
VIII. ξ-functions: Pólya's is massive and Riemann's is massless	14
IX. The massive Dirac Model in Rindler coordinates	16
X. The massless Dirac equation with delta function potentials	17
XI. Heuristic approach to the spectrum	20
XII. The Riemann zeros as spectrum and The Riemann hypothesis	23
1. Normalizable eigenstates	23
2. The Magnus expansion	24
3. The Mertens function and the Perron formula	25
XIII. The Riemann interferometer	28
XIV. Conclusions	29
Acknowledgements	30
References	30

I. INTRODUCTION

One of the most promising approaches to prove the Riemann Hypothesis [1]-[11] is based on the conjecture, due to Pólya and Hilbert, that the Riemann zeros are the eigenvalues of a quantum mechanical Hamiltonian [12]. This bold idea is supported by several results and analogies involving Number Theory, Random Matrix Theory and Quantum Chaos [13]-[29]. However the construction of a Hamiltonian whose spectrum contains the Riemann zeros, has eluded the researchers for several decades. In this paper we shall review the progress made along this direction starting from the famous xp model proposed in 1999 by Berry, Keating and Connes [30]-[32], that inspired many works [33]-[57], some of whom will be discussed below. See [58]-[62] for general reviews on physical approaches to the RH and [63]-[66] for introductions and historical background. Another approaches to the RH and related material can be found in [67]-[93].

To relate xp with the Riemann zeros, Berry, Keating and Connes used two different regularizations. The Berry and Keating regularization led to a discrete spectrum related to the smooth Riemann zeros [30, 31], while Connes's regularization led to an absorption spectrum where the *zeros* are missing spectral lines [32]. A physical realization of the Connes model was obtained in 2008 in terms of the dynamics of an electron moving in two dimensions under the action of a uniform perpendicular magnetic field and a electrostatic potential [41]. However this model has not been able to reproduce the exact location of the Riemann zeros. On the other hand, the Berry-Keating xp model, was revisited in 2011 in terms of the classical Hamiltonians $H = x(p+1/p)$, and $H = (x+1/x)(p+1/p)$ whose quantizations contain the smooth approximation of the Riemann zeros [45, 49]. Later on, these models were generalized in terms of the family of Hamiltonians $H = U(x)p + V(x)/p$, that were shown to describe the dynamics of a massive particle in a relativistic spacetime whose metric can be constructed using the functions U and V [48]. This result suggested a reformulation of $H = U(x)p + V(x)/p$ in terms of the massive Dirac equation in the aforementioned spacetimes [51]. Using this reformulation, the Hamiltonian $H = x(p+1/p)$ was shown to be equivalent to the massive Dirac equation in Rindler spacetime, that is the natural arena to study accelerated observers and the Unruh effect [55]. This result provides an appealing spacetime interpretation of the xp model and in particular of the smooth Riemann zeros.

To obtain the exact *zeros*, one has to make further modifications of the Dirac model. First of all, the fermion has to become massless. This change is suggested by a field theory interpretation of the Pólya's ξ function and its comparison with the Riemann's ξ function. On the other hand, inspired by the Berry's conjecture on the relation between prime numbers and periodic orbits [19, 22] we incorporated the prime numbers into the Dirac action by means of Dirac delta functions [55]. These delta functions represent moving mirrors that reflect or transmit massless fermions. The spectrum of the complete model can be analyzed using transfer matrix techniques that can be solved exactly in the limit where the reflection amplitudes of the mirrors go to zero, that is when the mirrors become transparent. In this limit we find that the *zeros* on the critical line are eigenvalues of the Hamiltonian by choosing appropriately the parameter that characterizes the self-adjoint extension of the Hamiltonian. One obtains in this manner a spectral realization of the Riemann zeros that differs from the Pólya and Hilbert conjecture in the sense that one needs to fine tune a parameter to *see* each individual *zero*. In our approach we are not able to find a single Hamiltonian encompassing all the *zeros* at once. Finally, we propose an experimental realization of the Riemann zeros using interferometer consisting in an array of semitransparent mirrors, or beam splitters, placed at positions related to the logarithms of the prime numbers.

The paper is organized in a historical and pedagogical way presenting at the end of each section a summary of achievements (✓), shortcomings/obstacles (✗) and questions/suggestions (?).

II. THE SEMICLASSICAL xp BERRY, KEATING AND CONNES MODEL

In this section we review the main results concerning the classical and semiclassical xp model [30-32]. A classical trajectory of the Hamiltonian $H = xp$, with energy E , is given by

$$x(t) = x_0 e^t, \quad p(t) = p_0 e^{-t}, \quad E = x_0 p_0, \quad (2.1)$$

that traces the parabola $E = xp$ in phase space plotted in Fig.1. E has the dimension of an action, so one should multiply xp by a frequency to get an energy, but for the time being we keep the notation $H = xp$. Under a time reversal transformation, $x \rightarrow x, p \rightarrow -p$ one finds $xp \rightarrow -xp$, so that this symmetry is broken. This is why reversing the time variable t in (2.1) does not yield a trajectory generated by xp . As $t \rightarrow \infty$, the trajectory becomes unbounded, that is $|x| \rightarrow \infty$, so one expects the semiclassical and quantum spectrum of the xp model to form a continuum. In order to get a discrete spectrum Berry and Keating introduced the constraints $|x| \geq \ell_x$ and $|p| \geq \ell_p$, so that the particle starts at $t = 0$ at $(x, p) = (\ell_x, E/\ell_x)$ and ends at $(x, p) = (E/\ell_p, \ell_p)$ after a time lapse $T = \log(E/\ell_x \ell_p)$ (we

assume for simplicity that $x, p > 0$). The trajectories are now bounded, but not periodic. A semiclassical estimate of the number of energy levels, $n_{\text{BK}}(E)$, between 0 and $E > 0$ is given by the formula

$$n_{\text{BK}}(E) = \frac{A_{\text{BK}}}{2\pi\hbar} = \frac{E}{2\pi\hbar} \left(\log \frac{E}{\ell_x \ell_p} - 1 \right) + \frac{7}{8}, \quad (2.2)$$

where A_{BK} is the phase space area below the parabola $E = xp$ and the lines $x = \ell_x$ and $p = \ell_p$, measured in units of the Planck's constant $2\pi\hbar$ (see Fig. 1). The term $7/8$ arises from the Maslow phase [30]. In the course of the paper, we shall encounter this equation several times with the constant term depending on the particular model.




FIG. 1: Left: The region in shadow describes the allowed phase space with area A_{BK} bounded by the classical trajectory (2.1) with $E > 0$ and the constraints $x \geq \ell_x, p \geq \ell_p$. Right: Same as before with the constraints $0 < x, p < \Lambda$.

Berry and Keating compared this result with the average number of Riemann zeros, whose imaginary part is less than t with $t \gg 1$,

$$\langle n(t) \rangle \simeq \frac{t}{2\pi} \left(\log \frac{t}{2\pi} - 1 \right) + \frac{7}{8} + O(1/t), \quad (2.3)$$

finding an agreement with the identifications

$$t = \frac{E}{\hbar}, \quad \ell_x \ell_p = 2\pi\hbar. \quad (2.4)$$

Thus, the semiclassical energies E , in units of \hbar , are identified with the Riemann zeros, while $\ell_x \ell_p$ is identified with the Planck's constant. This result is remarkable given the simplicity of the assumptions. But one must observe that the derivation of eq.(2.2) is heuristic, so one goal is to find a consistent quantum version of it.

Connes proposed another regularization of the xp model based on the restrictions $|x| \leq \Lambda$ and $|p| \leq \Lambda$, where Λ is a common cutoff, which is taken to infinity at the end of the calculation [32]. The semiclassical number of states is computed as before yielding (see Fig. 1, we set $\hbar = 1$)

$$n_{\text{C}}(E) = \frac{A_{\text{C}}}{2\pi} = \frac{E}{2\pi} \log \frac{\Lambda^2}{2\pi} - \frac{E}{2\pi} \left(\log \frac{E}{2\pi} - 1 \right). \quad (2.5)$$

The first term on the RHS of this formula diverges in the limit $\Lambda \rightarrow \infty$, which corresponds to a continuum of states. The second term is minus the average number of Riemann zeros, which according to Connes, become missing spectral lines in the continuum [11, 29, 32]. This is called the *absorption* spectral interpretation of the Riemann zeros, as opposed to the standard *emission* spectral interpretation where the *zeros* form a discrete spectrum. Connes, relates the minus sign in eq.(2.5) to a minus sign discrepancy between the fluctuation term of the number of zeros and the associated formula in the theory of Quantum Chaos. We shall show below that the negative term in Eq.(2.5) must be seen as a finite size correction of discrete energy levels and not as an indication of missing spectral lines.

Let us give for completeness the formula for the exact number of *zeros* up to t [2, 3]

$$\begin{aligned} n_{\mathbb{R}}(t) &= \langle n(t) \rangle + n_{\mathbb{H}}(t), \\ \langle n(t) \rangle &= \frac{\theta(t)}{\pi} + 1, \quad n_{\mathbb{H}}(t) = \frac{1}{\pi} \operatorname{Im} \log \zeta \left(\frac{1}{2} + it \right), \end{aligned} \quad (2.6)$$

where $\langle n(t) \rangle$ is the Riemann-von Mangoldt formula that gives the average behavior in terms of the function $\theta(t)$

$$\theta(t) = \operatorname{Im} \log \Gamma \left(\frac{1}{4} + \frac{it}{2} \right) - \frac{t}{2} \log \pi \xrightarrow{t \rightarrow \infty} \frac{t}{2} \log \frac{t}{2\pi} - \frac{t}{2} - \frac{\pi}{8} + O(1/t), \quad (2.7)$$

that can also be written as

$$e^{2i\theta(t)} = \pi^{-it} \frac{\Gamma \left(\frac{1}{4} + \frac{it}{2} \right)}{\Gamma \left(\frac{1}{4} - \frac{it}{2} \right)}. \quad (2.8)$$

$\theta(t)$ is the phase of the Riemann zeta function on the critical line, that can be expressed as

$$\zeta \left(\frac{1}{2} + it \right) = e^{-i\theta(t)} Z(t), \quad (2.9)$$

where $Z(t)$ is the Riemann-Siegel zeta function, or Hardy function, that on the critical line satisfies

$$Z(t) = Z(-t) = Z^*(t), \quad t \in \mathbb{R}. \quad (2.10)$$

Summary:

- ✓ The semiclassical xp Hamiltonian seems to be connected to the average Riemann zeros.
- ✗ There are two schemes leading to opposite physical realizations: emission vs absorption.
- ? Quantum version of the semiclassical xp models.

III. THE QUANTUM xp MODEL

To quantize the xp Hamiltonian, Berry and Keating used the normal ordered operator [30]

$$\hat{H} = \frac{1}{2} (x \hat{p} + \hat{p} x) = -i\hbar \left(x \frac{d}{dx} + \frac{1}{2} \right), \quad x \in \mathbb{R}, \quad (3.1)$$

where x belongs to the real line and $\hat{p} = -i\hbar d/dx$ is the momentum operator. If x is restricted to the positive half-line, then (3.1) is equivalent to the expression

$$\hat{H} = \sqrt{x} \hat{p} \sqrt{x}, \quad x \geq 0. \quad (3.2)$$

\hat{H} is an essentially self-adjoint operator acting on the Hilbert space $L^2(0, \infty)$ of square integrable functions in the half line $\mathbb{R}_+ = (0, \infty)$ [35, 36, 42]. The eigenfunctions, with eigenvalue E , are given by

$$\psi_E(x) = \frac{1}{\sqrt{2\pi\hbar}} x^{-\frac{1}{2} + \frac{iE}{\hbar}}, \quad x > 0, \quad E \in \mathbb{R}, \quad (3.3)$$

and the spectrum is the real line \mathbb{R} . The normalization of (3.3) is given by the Dirac's delta function

$$\langle \psi_E | \psi_{E'} \rangle = \int_0^\infty dx \psi_E^*(x) \psi_{E'}(x) = \delta(E - E'). \quad (3.4)$$

The eigenfunctions (3.3) form an orthonormal basis of $L^2(0, \infty)$, that is related to the Mellin transform in the same manner that the eigenfunctions of the momentum operator \hat{p} , on the real line, are related to the Fourier transform [36]. If one takes x in the whole real line, then the spectrum of the Hamiltonian (3.1) is doubly degenerate. This

degeneracy can be understood from the invariance of xp under the parity transformation $x \rightarrow -x, p \rightarrow -p$, which allows one to split the eigenfunctions with energy E into even and odd sectors

$$\psi_E^{(e)}(x) = \frac{1}{\sqrt{2\pi\hbar}} |x|^{-\frac{1}{2} + \frac{iE}{\hbar}}, \quad \psi_E^{(o)}(x) = \frac{\text{sign } x}{\sqrt{2\pi\hbar}} |x|^{-\frac{1}{2} + \frac{iE}{\hbar}}, \quad x \in \mathbb{R}, \quad E \in \mathbb{R}. \quad (3.5)$$

Berry and Keating computed the Fourier transform of the even wave function $\psi_E^{(e)}(x)$ [30]

$$\begin{aligned} \hat{\psi}_E^{(e)}(p) &= \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} dx \psi_E^{(e)}(x) e^{-ipx/\hbar} \\ &= \frac{1}{\sqrt{2\pi\hbar}} |p|^{-\frac{1}{2} - \frac{iE}{\hbar}} (2\hbar)^{iE/\hbar} \frac{\Gamma\left(\frac{1}{4} + \frac{iE}{2\hbar}\right)}{\Gamma\left(\frac{1}{4} - \frac{iE}{2\hbar}\right)}, \end{aligned} \quad (3.6)$$

which means that the position and momentum eigenfunctions are each other's time reversed, giving a physical interpretation of the phase $\theta(t)$, see Eq.(2.8). Choosing odd eigenfunctions leads to an equation similar to Eq.(3.6) in terms of the gamma functions $\Gamma\left(\frac{3}{4} \pm \frac{iE}{2\hbar}\right)$ that appear in the functional relation of the odd Dirichlet characters. Equation (3.6) is a consequence of the exchange $x \leftrightarrow p$ symmetry of the xp Hamiltonian, which seems to be an important ingredient of the xp model.

Comments:

- Removing Connes's cutoff, i.e. $\Lambda \rightarrow \infty$, gives the quantum Hamiltonians (3.1) or (3.2), whose spectrum is a continuum. This shows that the negative term in Eq.(2.5) does not correspond to missing spectral lines. In the next section we give a physical interpretation of this term in another context.
- xp is invariant under the scale transformation (dilations) $x \rightarrow Kx, p \rightarrow K^{-1}p$, with $K > 0$. An example of this transformation is the classical trajectory (2.1), whose infinitesimal generator is xp . Under dilations, $\ell_x \rightarrow K\ell_x, \ell_p \rightarrow K^{-1}\ell_p$, so the condition $\ell_x\ell_p = 2\pi\hbar$ is preserved. Berry and Keating suggested to use integer dilations $K = n$, corresponding to evolution times $\log n$, to write [30]

$$\psi_E(x) \rightarrow \sum_{n=1}^{\infty} \psi_E(nx) = \frac{1}{\sqrt{2\pi\hbar}} x^{-\frac{1}{2} + \frac{iE}{\hbar}} \sum_{n=1}^{\infty} \frac{1}{n^{\frac{1}{2} - \frac{iE}{\hbar}}} = \frac{1}{\sqrt{2\pi\hbar}} x^{-\frac{1}{2} + \frac{iE}{\hbar}} \zeta(1/2 - iE/\hbar). \quad (3.7)$$

If there exists a physical reason for this quantity to vanish one would obtain the Riemann zeros E_n . Equation (3.7) could be interpreted as the breaking of the continuous scale invariance to discrete scale invariance.

Summary:

- ✗ The normal order quantization of xp does not exhibit any trace of the Riemann zeros.
- ✓ The phase of the zeta function appears in the Fourier transform of the xp eigenfunctions.

IV. THE LANDAU MODEL AND XP

Let us consider a charged particle moving in a plane under the action of a perpendicular magnetic field and an electrostatic potential $V(x, y) \propto xy$ [41]. The Lagrangian describing the dynamics is given, in the Landau gauge, by

$$\mathcal{L} = \frac{\mu}{2}(\dot{x}^2 + \dot{y}^2) - \frac{eB}{c} \dot{y}x - e\lambda xy, \quad (4.1)$$

where μ is the mass, e the electric charge, B the magnetic field, c the speed of light and λ a coupling constant that parameterizes the electrostatic potential. There are two normal modes with real, ω_c , and imaginary, ω_h , angular frequencies, describing a cyclotronic and a hyperbolic motion respectively. In the limit where $\omega_c \gg |\omega_h|$, only the Lowest Landau Level (LLL) is relevant and the effective Lagrangian becomes

$$\mathcal{L}_{\text{eff}} = p\dot{x} - |\omega_h|xp, \quad p = \frac{\hbar y}{\ell^2}, \quad \ell = \left(\frac{\hbar c}{eB}\right)^{1/2}, \quad (4.2)$$

where ℓ is the magnetic length, which is proportional to the radius of the cyclotronic orbits in the LLL. The coordinates x and y , which commute in the 2D model, after the projection to the LLL become canonical conjugate variables, and the effective Hamiltonian is proportional to the xp Hamiltonian with the proportionality constant given by the angular frequency $|\omega_h|$ (this is the missing frequency factor mentioned in section II). The quantum Hamiltonian associated to the Lagrangian (4.1) is

$$\hat{H} = \frac{1}{2\mu} \left[\hat{p}_x + \left(\hat{p}_y + \frac{\hbar}{\ell^2} x \right)^2 \right] + e\lambda xy, \quad (4.3)$$

where $\hat{p}_x = -i\hbar\partial_x$ and $\hat{p}_y = -i\hbar\partial_y$. After a unitary transformation (4.3) becomes the sum of two commuting Hamiltonians corresponding to the cyclotronic and hyperbolic motions alluded to above

$$\begin{aligned} H &= H_c + H_h, \\ H_c &= \frac{\omega_c}{2}(\hat{p}^2 + q^2), \quad H_h = \frac{|\omega_h|}{2}(\hat{P}Q + Q\hat{P}). \end{aligned} \quad (4.4)$$

In the limit $\omega_c \gg |\omega_h|$ one has

$$\omega_c \simeq \frac{eB}{\mu c}, \quad |\omega_h| \sim \frac{\lambda c}{B}. \quad (4.5)$$

The unitary transformation that brings Eq. (4.3) into Eq.(4.4) corresponds to the classical canonical transformation

$$q = x + py, \quad p = p_x, \quad Q = -py, \quad P = y + p_x. \quad (4.6)$$

When $\omega_c \gg |\omega_h|$, the low energy states of H are the product of the lowest eigenstate of H_c , namely $\psi = e^{-q^2/2\ell^2}$, times the eigenstates of H_h that can be chosen as even or odd under the parity transformation $Q \rightarrow -Q$

$$\Phi_E^+(Q) = \frac{1}{|Q|^{\frac{1}{2}-iE}}, \quad \Phi_E^-(Q) = \frac{\text{sign}(Q)}{|Q|^{\frac{1}{2}-iE}}. \quad (4.7)$$

The corresponding wave functions are given by (we choose $|\omega_h| = 1$)

$$\psi_E^\pm(x, y) = C \int dQ e^{-iQy/\ell^2} e^{-(x-Q)^2/2\ell^2} \Phi_E^\pm(Q), \quad (4.8)$$

where C is a normalization constant, which yields

$$\begin{aligned} \psi_E^+(x, y) &= C_E^+ e^{-\frac{x^2}{2\ell^2}} M\left(\frac{1}{4} + \frac{iE}{2}, \frac{1}{2}, \frac{(x-iy)^2}{2\ell^2}\right), \\ \psi_E^-(x, y) &= C_E^-(x-iy) e^{-\frac{x^2}{2\ell^2}} M\left(\frac{3}{4} + \frac{iE}{2}, \frac{3}{2}, \frac{(x-iy)^2}{2\ell^2}\right), \end{aligned} \quad (4.9)$$

where $M(a, b, z)$ is a confluent hypergeometric function [94]. Fig.2 shows that the maximum of the absolute value of ψ_E^+ is attained on the classical trajectory $E = xy$ (in units of $\hbar = \ell = 1$). This 2D representation of the classical trajectories is possible because in the LLL x and y become canonical conjugate variables and consequently the 2D plane coincides with the phase space (x, p) .

To count the number of states with an energy below E one places the particle into a box: $|x| < L, |y| < L$ and impose the boundary conditions

$$\psi_E^+(x, L) = e^{ixL/\ell^2} \psi_E^+(L, x), \quad (4.10)$$

which identifies the outgoing particle at $x = L$ with the incoming particle at $y = L$ up to a phase. The asymptotic behavior $L \gg \ell$ of (4.9) is

$$\begin{aligned} \psi_E^+(L, x) &\simeq e^{-ixL/\ell^2 - x^2/2\ell^2} \frac{\Gamma(\frac{1}{2})}{\Gamma(\frac{1}{4} + \frac{iE}{2})} \left(\frac{L^2}{2\ell^2}\right)^{-\frac{1}{4} + \frac{iE}{2}}, \\ \psi_E^+(x, L) &\simeq e^{-x^2/2\ell^2} \frac{\Gamma(\frac{1}{2})}{\Gamma(\frac{1}{4} - \frac{iE}{2})} \left(\frac{L^2}{2\ell^2}\right)^{-\frac{1}{4} - \frac{iE}{2}}, \end{aligned} \quad (4.11)$$

that plugged into the BC (4.10) yields

$$\frac{\Gamma\left(\frac{1}{4} + \frac{iE}{2}\right)}{\Gamma\left(\frac{1}{4} - \frac{iE}{2}\right)} \left(\frac{L^2}{2\ell^2}\right)^{-iE} = 1, \quad (4.12)$$

or using Eq.(2.8)

$$e^{2i\theta(E)} \left(\frac{L^2}{2\pi\ell^2}\right)^{-iE} = 1. \quad (4.13)$$

Hence the number of states $n(E)$ with energy less than E is given by

$$n(E) \simeq \frac{E}{2\pi} \log\left(\frac{L^2}{2\pi\ell^2}\right) + 1 - \langle n(E) \rangle, \quad (4.14)$$

whose asymptotic behavior coincides with Connes's formula (2.5) for a cutoff $\Lambda = L/\ell$. In fact, the term $\langle n(E) \rangle$ is the exact Riemann-von Mangoldt formula (2.6).




FIG. 2: Plot of $|\psi_E^+(x, y)|$ for $E = 10$ in the region $-10 < x, y < 10$. Left: 3D representation, Right: density plot.

Summary:

- ✓ The Landau model with a xy potential provides a physical realization of Connes's xp model.
- ✓ The finite size effects in the spectrum are given by the Riemann-von Mangoldt formula.
- ✗ There are no missing spectral lines in the physical realizations of xp à la Connes.

V. THE xP MODEL REVISITED

An intuitive argument of why the quantum Hamiltonian $(x\hat{p} + \hat{p}x)/2$ has a continuum spectrum is that the classical trajectories of xp are unbounded. Therefore, to have a discrete spectrum one should modify xp in order to bound the trajectories. This is achieved by the classical Hamiltonian [45]

$$H_I = x \left(p + \frac{\ell_p^2}{p} \right), \quad x \geq \ell_x. \quad (5.1)$$

For $|p| \gg \ell_p$, a classical trajectory with energy E satisfies $E \simeq xp$, but for $|p| \sim \ell_p$, the coordinate $|x|$ slows down, reaches a maximum and goes back to the value ℓ_x , where it bounces off starting again at high momentum. In this manner one gets a periodic orbit (see fig. 3)




FIG. 3: Classical trajectories of the Hamiltonians (5.1) (left) and (5.4) (right) in phase space with $E > 0$. The dashed lines denote the hyperbola $E = xp$. (ℓ_x, ℓ_p) is a fixed point solution of the classical equations generated by (5.1) and (5.4).

$$\begin{aligned} x(t) &= \frac{\ell_x}{|p_0|} e^{2t} \sqrt{(p_0^2 + \ell_p^2) e^{-2t} - \ell_p^2}, \quad 0 \leq t \leq T_E, \\ p(t) &= \pm \sqrt{(p_0^2 + \ell_p^2) e^{-2t} - \ell_p^2}, \end{aligned} \quad (5.2)$$

where T_E is the period given by (we take $E > 0$)

$$T_E = \cosh^{-1} \frac{E}{2\ell_x \ell_p} \rightarrow \log \frac{E}{\ell_x \ell_p} \quad (E \gg \ell_x \ell_p). \quad (5.3)$$

The asymptotic value of T_E is the time lapse it takes a particle to go from $x = \ell_x$ to $x = E/\ell_p$ in the xp model.

The exchange symmetry $x \leftrightarrow p$ of xp is broken by the Hamiltonian (5.1). To restore it, Berry and Keating proposed the $x - p$ symmetric Hamiltonian [49]

$$H_{II} = \left(x + \frac{\ell_x^2}{x} \right) \left(p + \frac{\ell_p^2}{p} \right), \quad x \geq 0. \quad (5.4)$$

Here the classical trajectories turn clockwise around the point (ℓ_x, ℓ_p) , and for $x \gg \ell_x$ and $p \gg \ell_p$, approach the parabola $E = xp$ (see Fig. 3). The semiclassical analysis of (5.1) and (5.4) reproduce the asymptotic behavior of Eq.(2.2) to leading orders $E \log E$ and E , but differ in the remaining terms.

The two models discussed above have the general form

$$H = U(x)p + \ell_p^2 \frac{V(x)}{p}, \quad x \in D, \quad (5.5)$$

where $U(x)$ and $V(x)$ are positive functions defined in an interval D of the real line. H_I corresponds to $U(x) = V(x) = x, D = (\ell_x, \infty)$, and H_{II} corresponds to $U(x) = V(x) = x + \ell_x^2/x, D = (0, \infty)$. The classical Hamiltonian (5.5) can be quantized in terms of the operator

$$\hat{H} = \sqrt{U} \hat{p} \sqrt{U} + \ell_p^2 \sqrt{V} \hat{p}^{-1} \sqrt{V}, \quad (5.6)$$

where \hat{p}^{-1} is pseudo-differential operator

$$(\hat{p}^{-1} \psi)(x) = -\frac{i}{\hbar} \int_x^\infty dy \psi(y), \quad (5.7)$$




FIG. 4: Absolute values of the wave function $\psi_E(x)$, given in eq.(5.12) (continuous line), and $x^{-\frac{1}{2}}$ (dashed line).

which satisfies that $\hat{p}\hat{p}^{-1} = \hat{p}^{-1}\hat{p} = \mathbf{1}$ acting on functions which vanish in the limit $x \rightarrow \infty$. The action of \hat{H} is

$$(\hat{H}\psi)(x) = -i\hbar\sqrt{U(x)}\frac{d}{dx}\left\{\sqrt{U(x)}\psi(x)\right\} - \frac{i\ell_p^2}{\hbar}\int_x^\infty dy\sqrt{V(x)V(y)}\psi(y). \quad (5.8)$$

The normal order prescription that leads from (5.5) to (5.8) will be derived in section VII in the case where $U(x) = V(x) = x$, but holds in general [51]. We want the Hamiltonian (5.6) to be self-adjoint, that is [96, 97]

$$\langle\psi_1|\hat{H}|\psi_2\rangle = \langle\hat{H}\psi_1|\psi_2\rangle. \quad (5.9)$$

When the interval is $D = (\ell_x, \infty)$, Eq.(5.9) holds for wave functions that vanishes sufficiently fast at infinity and satisfy the non local boundary condition

$$\hbar e^{i\vartheta}\sqrt{U(\ell_x)}\psi(\ell_x) = \ell_p\int_{\ell_x}^\infty dx\sqrt{V(x)}\psi(x), \quad (5.10)$$

where $\vartheta \in [0, 2\pi)$ parameterizes the self-adjoint extensions of \hat{H} . The quantum Hamiltonian associated to (5.1) is

$$\hat{H}_I = \sqrt{x}\hat{p}\sqrt{x} + \ell_p^2\sqrt{x}\hat{p}^{-1}\sqrt{x}, \quad x \geq \ell_x, \quad (5.11)$$

and its eigenfunctions are proportional to (see fig. 4)

$$\psi_E(x) = x^{\frac{iE}{2\hbar}}K_{\frac{1}{2}-\frac{iE}{2\hbar}}\left(\frac{\ell_px}{\hbar}\right) \propto \begin{cases} x^{-\frac{1}{2}+\frac{iE}{\hbar}} & x \ll \frac{E}{2\ell_p}, \\ x^{-\frac{1}{2}+\frac{iE}{2\hbar}}e^{-\ell_px/\hbar} & x \gg \frac{E}{2\ell_p}, \end{cases} \quad (5.12)$$

where $K_\nu(z)$ is the modified K -Bessel function [94]. For small values of x , the wave functions (5.12) behave as those of the xp Hamiltonian, given in Eq.(3.3), while for large values of x they decay exponentially giving a normalizable state. The boundary condition (5.10) reads in this case

$$\hbar e^{i\vartheta}\sqrt{\ell_x}\psi(\ell_x) = \ell_p\int_{\ell_x}^\infty dx\sqrt{x}\psi(x), \quad (5.13)$$

and substituting (5.12) yields the equation for the eigenenergies E_n ,

$$e^{i\vartheta}K_{\frac{1}{2}-\frac{iE}{2\hbar}}\left(\frac{\ell_x\ell_p}{\hbar}\right) - K_{\frac{1}{2}+\frac{iE}{2\hbar}}\left(\frac{\ell_x\ell_p}{\hbar}\right) = 0. \quad (5.14)$$

For $\vartheta = 0$ or π , the eigenenergies form time reversed pairs $\{E_n, -E_n\}$, and for $\vartheta = 0$, there is a zero energy state $E = 0$. Considering that the Riemann zeros form pairs $s_n = 1/2 \pm it_n$, with t_n real under the RH, and that $s = 1/2$ is not a zero of $\zeta(s)$, we are led to the choice $\vartheta = \pi$. On the other hand, using the asymptotic behavior

$$K_{a+\frac{it}{2}}(z) \longrightarrow \sqrt{\frac{\pi}{t}}\left(\frac{t}{z}\right)^a e^{-\pi t/4} e^{\frac{i\pi}{2}(a-\frac{1}{2})}\left(\frac{t}{ze}\right)^{it/2}, \quad a > 0, t \gg 1, \quad (5.15)$$

one derives in the limit $|E| \gg \hbar$,

$$K_{\frac{1}{2} + \frac{iE}{2\hbar}} \left(\frac{\ell_x \ell_p}{\hbar} \right) + K_{\frac{1}{2} - \frac{iE}{2\hbar}} \left(\frac{\ell_x \ell_p}{\hbar} \right) = 0 \longrightarrow \cos \left(\frac{E}{2\hbar} \log \frac{E}{\ell_x \ell_p e} \right) = 0, \quad (5.16)$$

hence the number of eigenenergies in the interval $(0, E)$ is given asymptotically by

$$n(E) \simeq \frac{E}{2\pi\hbar} \left(\log \frac{E}{\ell_x \ell_p} - 1 \right) - \frac{1}{2} + O(E^{-1}). \quad (5.17)$$

This equation agrees with the leading terms of the semiclassical spectrum (2.2) and the average Riemann zeros (2.3) under the identifications (2.4). Concerning the classical Hamiltonian (5.4), Berry and Keating obtained, by a semiclassical analysis, the asymptotic behavior of the counting function $n(E)$

$$n(t) \simeq \frac{t}{2\pi} \left(\log \frac{t}{2\pi} - 1 \right) - \frac{8\pi}{t} \log \frac{t}{2\pi} + \dots, \quad t \gg 1, \quad (5.18)$$

where $t = E/\hbar$ and $\ell_x \ell_p = 2\pi\hbar$. Again, the first two leading terms agree with Riemann's formula (2.3), while the next leading corrections are different from (5.17). In both cases, the constant $7/8$ in Riemann's formula (2.3) is missing.

Summary:

- ✓ The Berry-Keating xp model can be implemented quantum mechanically.
- ✗ The classical xp Hamiltonian has to be modified with ad-hoc terms to have bounded trajectories.
- ✗ In the quantum theory the latter terms become non local operators.
- ✗ The modified xp quantum Hamiltonian related to the average Riemann zeros is not unique.
- ✗ There is no trace of the exact Riemann zeros in the spectrum of the modified xp models.

VI. THE SPACE-TIME GEOMETRY OF THE MODIFIED xp MODELS

In this section we show that the modified xp Hamiltonian (5.5) is a disguised general theory of relativity [48]. Let us first consider the Lagrangian of the xp model,

$$L = p\dot{x} - H = p\dot{x} - xp. \quad (6.1)$$

In classical mechanics, where $H = p^2/2m + V(x)$, the Lagrangian can be expressed solely in terms of the position x and velocity $\dot{x} = dx/dt$. This is achieved by writing the momentum in terms of the velocity by means of the Hamilton equation $\dot{x} = \partial H/\partial p = p/m$. However, in the xp model the momentum p is not a function of the velocity because $\dot{x} = \partial H/\partial p = x$. Hence the Lagrangian (6.1) cannot be expressed solely in terms of x and \dot{x} . The situation changes radically for the Hamiltonian (5.5) whose Lagrangian is given by

$$L = p\dot{x} - H = p\dot{x} - U(x)p - \ell_p^2 \frac{V(x)}{p}. \quad (6.2)$$

Here the equation of motion

$$\dot{x} = \frac{\partial H}{\partial p} = U(x) - \ell_p^2 \frac{V(x)}{p^2}, \quad (6.3)$$

allows one to write p in terms of x and \dot{x} ,

$$p = \eta \ell_p \sqrt{\frac{V(x)}{U(x) - \dot{x}}}, \quad \eta = \text{sign } p, \quad (6.4)$$

where $\eta = \pm 1$ is the sign of the momentum that is a conserved quantity. The positivity of $U(x)$ and $V(x)$, imply that the velocity \dot{x} must never exceed the value of $U(x)$. Substituting (6.4) back into (6.2), yields the action

$$S_\eta = -\ell_p \eta \int \sqrt{-ds^2}, \quad (6.5)$$




FIG. 5: Left: Domain \mathcal{U} of Minkowski spacetime given in Eq.(6.10). Right: The classical trajectory given in Eq.(5.2), and plotted in Fig.3-left, becomes a straightline that bounces off regularly at the boundary (dotted line).

which, for either sign of η , is the action of a relativistic particle moving in a 1+1 dimensional spacetime metric

$$ds^2 = 4V(x)(-U(x)dt^2 + dt dx). \quad (6.6)$$

The parameter ℓ_p plays the role of mc where m is the mass of the particle and c is the speed of light. This result implies that the classical trajectories of the Hamiltonian (5.5) are the geodesics of the metric (6.6). The unfamiliar form of (5.5) is due to a special choice of spacetime coordinates where the component g_{xx} of the metric vanishes. A diffeomorphism of x permits to set $V(x) = U(x)$. The scalar curvature of the metric (6.6), in this *gauge*, is

$$R(x) = -2 \frac{\partial_x^2 V(x)}{V(x)}, \quad (6.7)$$

and vanishes for the models $V(x) = x$ and $V(x) = \text{constant}$. For the Hamiltonian (5.4) one obtains $R(x) = -4\ell_x^2/(x(x^2 + \ell_x^2))$ which vanishes asymptotically.

The flatness of the metric associated to the Hamiltonian (5.1) implies the existence of coordinates x^0, x^1 where (6.6) takes the Minkowski form

$$ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu, \quad \text{diag } \eta_{\mu\nu} = (-1, 1). \quad (6.8)$$

The change of variables is given by

$$t = \frac{1}{2} \log(x^0 + x^1), \quad x = \sqrt{-(x^0)^2 + (x^1)^2}. \quad (6.9)$$

Let \mathcal{U} denote the space-time domain of the model. In both coordinates it reads

$$\mathcal{U} = \{(t, x) \mid t \in (-\infty, \infty), x \geq \ell_x\} = \{(x^0, x^1) \mid x^0 \in (-\infty, \infty), x^1 \geq \sqrt{(x^0)^2 + \ell_x^2}\}. \quad (6.10)$$

The boundary of \mathcal{U} , denoted by $\partial\mathcal{U}$, is the hyperbola $x^1 = \sqrt{(x^0)^2 + \ell_x^2}$, that passes through the point $(x^0, x^1) = (0, \ell_x)$, (see fig 5). A convenient parametrization of the coordinates x^μ is given by the Rindler variables ρ and ϕ [98]

$$x^0 = \rho \sinh \phi, \quad x^1 = \rho \cosh \phi, \quad (6.11)$$

or in light-cone coordinates

$$x^\pm = x^0 \pm x^1 = \pm \rho e^{\pm \phi}, \quad (6.12)$$

where the Minkowski metric becomes

$$ds^2 = -dx^+ dx^- = d\rho^2 - \rho^2 d\phi^2. \quad (6.13)$$

These coordinates describe the right wedge of Rindler spacetime in 1+1 dimensions

$$\mathcal{R}_+ = \{(x^0, x^1) \mid x^0 \in (-\infty, \infty), x^1 \geq |x^0|\} = \{(\rho, \phi) \mid \phi \in (-\infty, \infty), \rho > 0\}. \quad (6.14)$$

Notice that $\mathcal{U} \subset \mathcal{R}_+$. The boundary $\partial\mathcal{U}$ corresponds to the hyperbola $\rho = \ell_x$ that is the worldline of a particle moving with uniform acceleration equal to $1/\ell_x$ (in units $c = 1$). The Rindler variables are the ones used to study the Unruh effect [95].

Let us now consider the classical Hamiltonian (5.4). The underlying metric is given by Eq.(6.6) with $U(x) = V(x) = x + \ell_x^2/x$. The change of variables

$$t = \frac{1}{2} \log(x^0 + x^1), \quad x = \sqrt{-(x^0)^2 + (x^1)^2 - \ell_x^2}, \quad (6.15)$$

brings the metric to the form

$$ds^2 = \frac{-(x^0)^2 + (x^1)^2}{-(x^0)^2 + (x^1)^2 - \ell_x^2} \eta_{\mu\nu} dx^\mu dx^\nu = \frac{\rho^2}{\rho^2 - \ell_x^2} (d\rho^2 - \rho^2 d\phi^2), \quad \rho \geq \ell_x, \quad (6.16)$$

which in the limit $\rho \rightarrow \infty$ converges to the flat metric (6.13).

Summary:

- ✓ The classical modified xp models are general relativistic theories in 1+1 dimensions.
- ✓ $H = x(p + \ell_p^2/p)$ is related to a domain \mathcal{U} of Rindler space-time.
- ✓ l_p is the mass of the particle.
- ✓ $1/\ell_x$ is the acceleration of a particle whose worldline is the boundary of \mathcal{U} .
- ? Relativistic quantum field theory of the modified xp models.

VII. DIRACIZATION OF $H = X(P + \ell_p^2/P)$

In this section we show that the Dirac theory provides the relativistic quantum version of the modified xp models [55]. We shall focus on the classical Hamiltonian $H = x(p + \ell_p^2/p)$ because the flatness of the associated space-time makes the computations easier, but the result is general: the quantum Hamiltonian (5.8) can be derived from the Dirac equation in a curved space-time with metric (6.6) [51].

The Dirac action of a fermion with mass m in the spacetime domain (6.10) is given by (in units $\hbar = c = 1$)

$$S = \frac{i}{2} \int_{\mathcal{U}} dx^0 dx^1 \bar{\psi} (\not{\partial} + im) \psi, \quad (7.1)$$

where ψ is a two component spinor, $\bar{\psi} = \psi^\dagger \gamma^0$, $\not{\partial} = \gamma^\mu \partial_\mu$ ($\partial_\mu = \partial/\partial x^\mu$), and γ^μ are the 2d Dirac matrices written in terms of the Pauli matrices $\sigma^{x,y}$ as

$$\gamma^0 = \sigma^x, \quad \gamma^1 = -i\sigma^y, \quad \psi = \begin{pmatrix} \psi_- \\ \psi_+ \end{pmatrix}. \quad (7.2)$$

The variational principle applied to (7.1) provides the Dirac equation

$$(\not{\partial} + im)\psi = 0, \quad (7.3)$$

and the boundary condition

$$\dot{x}^- \psi_-^\dagger \delta\psi_- - \dot{x}^+ \psi_+^\dagger \delta\psi_+ = 0, \quad (7.4)$$

where $\dot{x}^\pm = dx^\pm/d\phi = \ell_x e^{\pm\phi}$ is the vector tangent to the boundary $\partial\mathcal{U}$ in the light-cone coordinates $x^\pm = x^0 \pm x^1$. The Dirac equation reads in components

$$(\partial_0 - \partial_1)\psi_+ + im\psi_- = 0, \quad (\partial_0 + \partial_1)\psi_- + im\psi_+ = 0. \quad (7.5)$$

If $m = 0$ then ψ_{\pm} depends only x^{\pm} , and so the fields propagate to the left, $\psi_+(x^+)$, or to the right, $\psi_-(x^-)$, at the speed of light. The derivatives in Eq.(7.5) can be written in terms the variables t and x using Eq.(6.9),

$$\partial_0 - \partial_1 = -\frac{2e^{2t}}{x}\partial_x, \quad \partial_0 + \partial_1 = e^{-2t}(\partial_t + x\partial_x). \quad (7.6)$$

Let us denote by $\tilde{\psi}_{\mp}(t, x)$ the fermion fields in the coordinates t, x and by $\psi_{\mp}(x^0, x^1)$ the fields in the coordinates x^0, x^1 . The relation between these fields is given by the transformation law

$$\psi_- = \left(\frac{\partial x}{\partial x^-}\right)^{\frac{1}{2}} \tilde{\psi}_- = (2x)^{-\frac{1}{2}} e^t \tilde{\psi}_-, \quad \psi_+ = \left(\frac{\partial x}{\partial x^+}\right)^{\frac{1}{2}} \tilde{\psi}_- = (x/2)^{\frac{1}{2}} e^{-t} \tilde{\psi}_+. \quad (7.7)$$

Plugging Eqs.(7.6) and (7.7) into (7.5) gives

$$i\partial_t \tilde{\psi}_- = -i\sqrt{x}\partial_x (\sqrt{x}\tilde{\psi}_-) + m x \tilde{\psi}_+, \quad \partial_x (\sqrt{x}\tilde{\psi}_+) = im\sqrt{x}\tilde{\psi}_-. \quad (7.8)$$

The second equation is readily integrated

$$\tilde{\psi}_+(x, t) = -\frac{im}{\sqrt{x}} \int_x^{\infty} dy \sqrt{y} \tilde{\psi}_-(y, t), \quad (7.9)$$

and replacing it into the first equation in (7.8) gives

$$i\partial_t \tilde{\psi}_-(x, t) = -i\sqrt{x}\partial_x (\sqrt{x}\tilde{\psi}_-) - im^2\sqrt{x} \int_x^{\infty} dy \sqrt{y} \tilde{\psi}_-(y, t). \quad (7.10)$$

This is the Schroedinger equation with Hamiltonian (5.11) and the relation $m = \ell_p$ found in the previous section. The non locality of the Hamiltonian (5.11) is a consequence of the special coordinates t, x where the component $\tilde{\psi}_+$ becomes non dynamical and depends non locally on the component $\tilde{\psi}_-$ that is identified with the wave function of the modified xp model. Similarly, the boundary condition (5.13) can be derived from the Eq.(7.4) as follows. In Rindler coordinates the latter equation reads

$$e^{-\phi} \psi_-^{\dagger}(\ell_x, \phi) \delta \psi_-(\ell_x, \phi) = e^{\phi} \psi_+^{\dagger}(\ell_x, \phi) \delta \psi_+(\ell_x, \phi), \quad \forall \phi, \quad (7.11)$$

that is solved by

$$-ie^{i\vartheta} e^{-\phi/2} \psi_-(\ell_x, \phi) = e^{\phi/2} \psi_+(\ell_x, \phi), \quad \forall \phi, \quad (7.12)$$

where $\vartheta \in [0, 2\pi)$. Using Eq.(7.7) this equation becomes

$$-ie^{i\vartheta} \tilde{\psi}_-(\ell_x, t) = \tilde{\psi}_+(\ell_x, t), \quad \forall t, \quad (7.13)$$

that together with Eq.(7.9) yields Eq.(5.13). This completes the derivation of the quantum Hamiltonian and boundary condition associated to $H = x(p + \ell_p^2/p)$. The eigenfunctions and eigenvalue equation of this model were found in Section V. However, we shall rederive them in alternative way that will provide new insights in the next section.

Let us start by constructing the plane wave solutions of the Dirac equation (7.5),

$$\begin{pmatrix} \psi_- \\ \psi_+ \end{pmatrix} \propto \begin{pmatrix} e^{i\pi/4} e^{\beta/2} \\ e^{-i\pi/4} e^{-\beta/2} \end{pmatrix} e^{i(-p^0 x^0 + p^1 x^1)}, \quad (7.14)$$

where (p^0, p^1) is the energy-momentum vector parameterized in terms of the rapidity variable β

$$\begin{aligned} (p^0)^2 - (p^1)^2 &= m^2, \\ p^0 &= im \sinh \beta, \quad p^1 = im \cosh \beta, \quad \beta \in (-\infty, \infty). \end{aligned} \quad (7.15)$$

In Rindler coordinates these plane wave solutions decay exponentially with the distance as corresponds to a localized wave function

$$e^{i(-p^0 x^0 + p^1 x^1)} = e^{-m\rho \cosh(\beta - \phi)} \rightarrow 0, \quad \text{as } \rho \rightarrow \infty. \quad (7.16)$$

The general solution of the Dirac equation is given by the linear superposition of plane waves (7.14). The superposition that reproduces the eigenfunctions of the modified xp model is

$$\begin{aligned}\psi_{\mp}(\rho, \phi) &= e^{\pm i\pi/4} \int_{-\infty}^{\infty} d\beta e^{-iE\beta/2} e^{\pm\beta/2} e^{-m\rho \cosh(\beta-\phi)} \\ &= 2e^{\pm i\pi/4} e^{(\pm\frac{1}{2} - \frac{iE}{2})\phi} K_{\frac{1}{2} \mp \frac{iE}{2}}(m\rho),\end{aligned}\quad (7.17)$$

that replaced in Eq.(7.12) gives

$$e^{i\vartheta} K_{\frac{1}{2} - \frac{iE}{2}}(m\ell_x) - K_{\frac{1}{2} + \frac{iE}{2}}(m\ell_x) = 0, \quad (7.18)$$

which coincides with the eigenvalue equation (5.14) with $m = \ell_p$. Setting $m\ell_x = 2\pi$ and $\vartheta = \pi$, brings Eq.(7.18) to the form

$$\xi_H(t) \equiv K_{\frac{1}{2} + \frac{it}{2}}(2\pi) + K_{\frac{1}{2} - \frac{it}{2}}(2\pi) = 0. \quad (7.19)$$

Summary:

- ✓ The spectrum of a relativistic massive fermion in the domain \mathcal{U} agrees with the average Riemann zeros.
 ? Does this result provide a hint on a physical realization of the Riemann zeros.

VIII. ξ -FUNCTIONS: PÓLYA'S IS MASSIVE AND RIEMANN'S IS MASSLESS

The function $\xi_H(t)$ appearing in Eq.(7.19) reminds the *fake* ξ function defined by Pólya in 1926 [99, 100]

$$\xi^*(t) = 4\pi^2 \left(K_{\frac{9}{4} + \frac{it}{2}}(2\pi) + K_{\frac{9}{4} - \frac{it}{2}}(2\pi) \right). \quad (8.1)$$

This function shares several properties with the Riemann ξ function

$$\xi(t) = \frac{1}{4} s(s-1) \Gamma\left(\frac{s}{2}\right) \pi^{-s/2} \zeta(s), \quad s = \frac{1}{2} + it, \quad (8.2)$$

namely, $\xi^*(t)$ is an entire and even function of t , its zeros lie on the real axis and behave asymptotically like the average Riemann zeros, as shown by the expansion obtained using Eq.(5.15)

$$\xi^*(t) \xrightarrow{t \rightarrow \infty} 2^{3/4} \pi^{-7/4} t^{7/4} e^{-\pi t/4} \cos\left(\frac{t}{2} \log\left(\frac{t}{2\pi e}\right) + \frac{7\pi}{8}\right). \quad (8.3)$$

The zeros of $\xi(t)$, $\xi_H(t)$ and $\xi^*(t)$ are plotted in fig.6. The slight displacement between the two top curves is due to the constant $7\pi/8$ appearing in the argument of the cosine function in Eq.(8.3) as compared to that in Eq.(5.16).




FIG. 6: From bottom to top: plot of $-\log|\xi(t)|$ (Riemann zeros), $-\log|\xi_H(t)|$ (eigenvalues of the Hamiltonian (5.11) with $\ell_x \ell_p = 2\pi$) and $-\log|\xi^*(t)|$ (Pólya zeros). The cusp represents the zeros of the corresponding functions.

The similarity between $\xi_H(t)$ and $\xi^*(t)$, and the relation between $\xi^*(t)$ and $\xi(t)$ provide a hint on the field theory underlying the Riemann zeros. To show this, we shall review how Pólya arrived at $\xi^*(t)$. The starting point is the expression of $\xi(t)$ as a Fourier transform [3]

$$\begin{aligned}\xi(t) &= 4 \int_1^\infty dx \frac{d[x^{\frac{3}{2}}\psi'(x)]}{dx} x^{-\frac{1}{4}} \cos\left(\frac{t \log x}{2}\right), \\ \psi(x) &= \sum_{n=1}^\infty e^{-n^2 \pi x}, \quad \psi'(x) = \frac{d\psi(x)}{dx}.\end{aligned}\tag{8.4}$$

In the variable $x = e^\beta$ these equations become,

$$\begin{aligned}\xi(t) &= \int_0^\infty d\beta \Phi(\beta) \cos \frac{t\beta}{2}, \\ \Phi(\beta) &= 2\pi e^{5\beta/4} \sum_{n=1}^\infty (2\pi e^\beta n^2 - 3) n^2 e^{-\pi n^2 e^\beta}.\end{aligned}\tag{8.5}$$

The function $\Phi(\beta)$ behaves asymptotically as

$$\Phi(\beta) \rightarrow 4\pi^2 e^{9\beta/4} e^{-\pi e^\beta}, \quad \beta \rightarrow \infty,\tag{8.6}$$

which Pólya replaced by the following expression (see Fig. 7).

$$\Phi^*(\beta) = 4\pi^2 \left(e^{9\beta/4} + e^{-9\beta/4} \right) e^{-\pi(e^\beta + e^{-\beta})}.\tag{8.7}$$

The function $\xi^*(t)$ is defined as the Fourier transform of $\Phi^*(\beta)$,




FIG. 7: Plot of $\Phi(\beta)$ (red on line), and $\Phi^*(\beta)$ (blue on line). Outside the region $|\beta| < 1$ the difference is very small.

$$\xi^*(t) = \int_0^\infty d\beta \Phi^*(\beta) \cos \frac{t\beta}{2},\tag{8.8}$$

which gives finally the Eq.(8.1). The function (7.19) can also be written as the Fourier transform

$$\xi_H(t) = \int_0^\infty d\beta \Phi_H(\beta) \cos \frac{t\beta}{2},\tag{8.9}$$

with

$$\Phi_H(\beta) = (e^{\beta/2} + e^{-\beta/2}) e^{-2\pi \cosh \beta}\tag{8.10}$$

Observe that the term $e^{-2\pi \cosh \beta}$ appears in $\Phi_H(\beta)$ and $\Phi^*(\beta)$. The origin of this term in the Dirac theory is the plane wave factor (7.16) of a fermion with mass m located at the boundary $\rho = \ell_x$ with $m\ell_x = 2\pi$. This observation

suggests that the Pólya ξ function arises in the relativistic theory of a massive particle with scaling dimension $9/4$, rather than $1/2$, that corresponds to a fermion (this would explain the different order of the corresponding Bessel functions). The approximation $\Phi(\beta) \simeq \Phi^*(\beta)$, that is $e^{-\pi e^\beta} \simeq e^{-2\pi \cosh \beta}$, can then be understood as the replacement of a massless particle by a massive one. Indeed, the energy-momentum of a massless right moving particle is given by $p^0 = p^1 = \Lambda e^\beta$, where Λ is an energy scale. The corresponding plane wave factor is $e^{-\pi e^\beta}$, with $\Lambda = \pi$. For large rapidities, $\beta \gg 1$, a massive particle behaves as a massless one, i.e. $e^{-2\pi \cosh \beta} \simeq e^{-\pi e^\beta}$. However, for small rapidities this is not the case. These arguments suggest that the field theory underlying the Riemann ξ function, if it exists, must associated to a massless particle.

Summary:

- ✓ The zeros of the Polya ξ^* function behave as the spectrum of a relativistic massive particle in the domain \mathcal{U} .
- ? Polya's construction of ξ^* suggests that the Riemann's ξ function is related to a massless particle.

IX. THE MASSIVE DIRAC MODEL IN RINDLER COORDINATES

Let us formulate the Dirac theory in Rindler coordinates. Under a Lorentz transformation with boost parameter λ , the light cone coordinates x^\pm and the Dirac spinors ψ_\pm transform as

$$x^\pm \rightarrow e^{\mp\lambda} x^\pm, \quad \psi_\pm \rightarrow e^{\pm\lambda/2} \psi_\pm, \quad (9.1)$$

and the Rindler coordinates (6.11) as

$$\phi \rightarrow \phi - \lambda, \quad \rho \rightarrow \rho. \quad (9.2)$$

Hence the new spinor fields χ_\pm defined as

$$\chi_\pm = e^{\pm\phi/2} \psi_\pm, \quad (9.3)$$

remain invariant under (9.2). The Rindler wedge \mathcal{R}_+ , and its domain \mathcal{U} , are also invariant under Lorentz transformations. The Dirac action (7.1) written in terms of the spinors χ_\pm reads

$$S = \frac{i}{2} \int_{-\infty}^{\infty} d\phi \int_{\ell_x}^{\infty} d\rho \left[\chi_-^\dagger (\partial_\phi + \rho \partial_\rho + \frac{1}{2}) \chi_- + \chi_+^\dagger (\partial_\phi - \rho \partial_\rho - \frac{1}{2}) \chi_+ + im\rho (\chi_-^\dagger \chi_+ + \chi_+^\dagger \chi_-) \right], \quad (9.4)$$

while the Dirac equation (7.3) and the boundary condition (7.12) become

$$(\partial_\phi \pm \rho \partial_\rho \pm \frac{1}{2}) \chi_\mp + im\rho \chi_\pm = 0, \quad (9.5)$$

and

$$-ie^{i\vartheta} \chi_- = \chi_+ \quad \text{at } \rho = \ell_x. \quad (9.6)$$

The infinitesimal generator of translations of the Rindler time ϕ , acting on the spinor wave functions, is the Rindler Hamiltonian H_R , that can be read off from (9.5)

$$i\partial_\phi \chi = H_R \chi, \quad \chi = \begin{pmatrix} \chi_- \\ \chi_+ \end{pmatrix}, \quad (9.7)$$

$$H_R = \begin{pmatrix} -i(\rho \partial_\rho + \frac{1}{2}) & m\rho \\ m\rho & i(\rho \partial_\rho + \frac{1}{2}) \end{pmatrix} = \sqrt{\rho} \hat{p}_\rho \sqrt{\rho} \sigma^z + m\rho \sigma^x, \quad (9.8)$$

where $\hat{p}_\rho = -i\partial/\partial\rho$, is the momentum operator conjugate to the radial coordinate ρ . Notice that the operator

$$H_{\rho p_\rho} = -i(\rho \partial_\rho + \frac{1}{2}) = \frac{1}{2}(\rho \hat{p}_\rho + \hat{p}_\rho \rho) = \sqrt{\rho} \hat{p}_\rho \sqrt{\rho}, \quad (9.9)$$

coincides with Eq.(3.1) with the identification $x = \rho$ (in units $\hbar = 1$). The eigenfunctions of (9.9) are

$$H_{\rho p_\rho} \psi_E = E \psi_E, \quad \psi_E = \frac{1}{\sqrt{2\pi}} \rho^{-1/2+iE}, \quad (9.10)$$

with real eigenvalue E for $\rho > 0$ (recall Eq.(3.3)). Thus H_R consists of two copies of xp , with different signs corresponding to opposite fermion chiralities that are coupled by the mass term $m\rho\sigma^x$.

The scalar product of two wave functions, in the domain \mathcal{U} , can be defined as

$$\langle \chi_1 | \chi_2 \rangle = \int_{\ell_x}^{\infty} d\rho (\chi_{1,-}^* \chi_{2,-} + \chi_{1,+}^* \chi_{2,+}). \quad (9.11)$$

The Hamiltonian H_R is hermitean with this scalar product acting on wave functions that satisfy Eq.(9.6) and vanish sufficiently fast at infinity, i.e. $\lim_{\rho \rightarrow \infty} \rho^{1/2} \chi_{\pm}(\rho, \phi) = 0$. The eigenvalues and eigenvectors of the Hamiltonian (9.8), are given by the solutions of the Schroedinger equation

$$H_R \chi = E_R \chi, \quad \chi_{\pm}(\rho, \phi) = e^{-iE_R \phi \mp i\pi/4} K_{\frac{1}{2} \pm iE_R}(m\rho), \quad \rho \geq \ell_x, \quad (9.12)$$

which coincide with Eq.(7.17) with the identification

$$E_R = \frac{E}{2}. \quad (9.13)$$

The factor of 1/2 comes from the relation $e^{2t} = x^0 + x^1 = \rho e^{\phi}$ (see Eq.(6.9)), that implies $e^{-iE_R \phi} \propto e^{-iEt}$. The Rindler eigenenergies are obtained replacing E by $2E_R$ in Eq.(7.18).

Comments:

- The Dirac Hamiltonian associated to the metric (6.16) is

$$H = \begin{pmatrix} h & m\rho\Lambda \\ m\rho\Lambda & -h \end{pmatrix}, \quad h = -i \left(\rho \partial_\rho + \frac{1}{2} + \frac{1}{2} \rho \partial_\rho (\log \Lambda) \right), \quad \Lambda = \frac{\rho}{\sqrt{\rho^2 - \ell_x^2}}. \quad (9.14)$$

In the limit $\rho \gg \ell_x$ this Hamiltonian converges towards (9.8).

- Gupta, Harikumar and de Queiroz proposed the Hamiltonian $(xp + \not{x})/2$ as a Dirac variant of the xp Hamiltonian [50]. The Hamiltonian is defined on a semi-infinite cylinder and becomes effectively one dimensional by considering the winding modes on the compact dimension. The eigenfunctions are given by Whittaker functions and the spectrum satisfies an equation similar to Eq.(4.12) in the Landau theory. In the limit where a regularization parameter goes to zero one obtains a continuum spectrum with a correction term related to the Riemann-von Mangoldt formula.

Summary:

- ✓ The massless Dirac Hamiltonian in Rindler spacetime is the direct sum of xp and $-xp$.
- ✓ The mass term couples the left and right modes of the fermions.

X. THE MASSLESS DIRAC EQUATION WITH DELTA FUNCTION POTENTIALS

From analogies between the Polya ξ^* function, the Riemann ξ function and the ξ_H function of the massive Dirac model, we conjectured in section VIII the existence of a massless field theory underlying ξ . At first look this idea does not look correct because the Hamiltonian obtained by setting $m = 0$ in Eq.(9.8), is equivalent to two copies of the quantum xp model which has a continuum spectrum. In fact, the mass term in that Hamiltonian is the mechanism responsible for obtaining a discrete spectrum.




FIG. 8: Left: worldlines of the mirrors with accelerations $a_n = 1/\ell_n = 1/n$ ($n = 1, 2, \dots$). Right: A massless fermion (dotted line) at the point $(x^0, x^1) = (0, 1)$ moves to the right until it hits a moving mirror where it can be reflected or transmitted.

To resolve this puzzle we shall replace the *bulk* mass term in the Dirac action (9.4) by a sum of ultra-local interactions placed at fixed values ℓ_n of the radial coordinate ρ [55]. These interactions can arise from moving mirrors, or beam splitters, that move with a uniform acceleration $1/\ell_n$ (see Fig. 8). The fermion moves freely, until it hits one of the mirrors and it is reflected or transmitted. The moving mirrors are realized mathematically by delta functions added to the massless Dirac action that couple the left and right components of the fermion on both sides of the mirror. These delta functions provide the matching conditions for the wave functions and can be parameterized by a complex number ϱ_n with $n = 2, \dots, \infty$. The scattering of the fermion at each mirror preserves unitarity that is equivalent to the self-adjointness of the Hamiltonian.

The model is formulated in the spacetime \mathcal{U} defined in Eq.(6.10). We divide \mathcal{U} into an infinite number of domains separated by hyperbolas with constant values of $\rho = \ell_n$, as follows. First we define the intervals (see Fig.9)

$$I_n = \{\rho \mid \ell_n < \rho < \ell_{n+1}\}, \quad n = 1, 2, \dots, \infty, \quad (10.1)$$

where using the scale invariance of the model we set $\ell_1 = 1$ (ℓ_1 plays the role of ℓ_x in previous sections).




FIG. 9: Intervals I_n defined in Eq.(10.1)

The partition of \mathcal{U} is given by

$$\mathcal{U} \rightarrow \tilde{\mathcal{U}} = \cup_{n=1}^{\infty} \mathcal{U}_n, \quad \mathcal{U}_n = \mathcal{I}_n \times \mathbb{R}, \quad (10.2)$$

where the factor \mathbb{R} denotes the range of the Rindler time ϕ . See Fig. 8 for an example with $\ell_n = n$. The wave function of the model is the two component Dirac spinor (see Eq.(9.7))

$$\chi(\rho) = \begin{pmatrix} \chi_-(\rho) \\ \chi_+(\rho) \end{pmatrix}, \quad \rho \in \mathcal{I} = \cup_{n=1}^{\infty} I_n, \quad (10.3)$$

and the scalar product is given by (recall Eq.(9.11))

$$\langle \chi | \chi \rangle = \sum_{n=1}^{\infty} \int_{\ell_n}^{\ell_{n+1}} d\rho \chi^\dagger(\rho) \cdot \chi(\rho). \quad (10.4)$$

The complex Hilbert space is $\mathcal{H} = L^2(\mathcal{I}, \mathbb{C}) \oplus L^2(\mathcal{I}, \mathbb{C})$ and the Hamiltonian is obtained setting $m = 0$ in Eq.(9.8)

$$H = \begin{pmatrix} -i(\rho \partial_\rho + \frac{1}{2}) & 0 \\ 0 & i(\rho \partial_\rho + \frac{1}{2}) \end{pmatrix}, \quad \rho \notin \mathcal{I}. \quad (10.5)$$

H is a self-adjoint operator acting on the subspace $\mathcal{H}_\vartheta \subset \mathcal{H}$ of wave functions that satisfy the boundary conditions [55] (see [101] for the relation between self-adjointness of operators and boundary conditions)

$$\chi \in \mathcal{H}_\vartheta : \quad \chi(\ell_n^-) = L(\varrho_n) \chi(\ell_n^+), \quad (n \geq 2), \quad -ie^{i\vartheta} \chi_-(\ell_1^+) = \chi_+(\ell_1^+), \quad (10.6)$$

where

$$\chi(\ell_n^\pm) = \lim_{\varepsilon \rightarrow 0^+} \chi(\ell_n \pm \varepsilon), \quad (10.7)$$

and

$$\vartheta \in [0, 2\pi), \quad L(\varrho) = \frac{1}{1 - |\varrho|^2} \begin{pmatrix} 1 + |\varrho|^2 & 2i\varrho \\ -2i\varrho^* & 1 + |\varrho|^2 \end{pmatrix}, \quad \varrho \in \mathbb{C}, \quad |\varrho| \neq 1. \quad (10.8)$$

This means that H satisfies

$$\langle \chi_1 | H \chi_2 \rangle = \langle H \chi_1 | \chi_2 \rangle, \quad \chi_{1,2} \in \mathcal{H}_\vartheta. \quad (10.9)$$

This condition guarantees that the norm (10.4) of the state is conserved by the time evolution generated by the Hamiltonian. The subspace \mathcal{H}_ϑ also depends on ℓ_n and ϱ_n but we shall not write this dependence explicitly. Similarly, we shall also denote the Hamiltonian as H_ϑ . The matching conditions (10.6) describe a scattering process where two incoming waves χ_n^{in} collide at the n^{th} -mirror and become two outgoing waves χ_n^{out} given by (see Fig. 10)

$$\chi_n^{\text{in}} = \begin{pmatrix} \chi_-(\ell_n^-) \\ \chi_+(\ell_n^+) \end{pmatrix}, \quad \chi_n^{\text{out}} = \begin{pmatrix} \chi_-(\ell_n^+) \\ \chi_+(\ell_n^-) \end{pmatrix}, \quad n > 1. \quad (10.10)$$

At the mirror $n = 1$, the components $\chi_\pm(\ell_1^-)$ of these vectors are null, i.e. there is propagation at the left of the boundary. The scattering process is described by the matrix S_n

$$\chi_n^{\text{out}} = S_n \chi_n^{\text{in}}, \quad S_n = \frac{1}{1 + |\varrho_n|^2} \begin{pmatrix} 1 - |\varrho_n|^2 & -2i\varrho_n \\ -2i\varrho_n^* & 1 - |\varrho_n|^2 \end{pmatrix}, \quad n > 1, \quad (10.11)$$

that is unitary,

$$S_n S_n^\dagger = \mathbf{1}. \quad (10.12)$$

Notice that the boundary condition at $\rho = \ell_1$, is also described by Eq.(10.11) with a parameter ϱ_1

$$\varrho_1 = -e^{-i\vartheta}, \quad (10.13)$$

that is a pure phase for the Hamiltonian H_ϑ to be self-adjoint. The matrix $L(\varrho)$ satisfies

$$L(1/\varrho^*) = -L(\varrho). \quad (10.14)$$

Hence, replacing ϱ_n by $1/\varrho_n^*$ gives a unitary equivalent model because the sign change at $\rho = \ell_n$, given in Eq.(10.14), can be compensated by changing the sign of the wave function in the remaining intervals. Hence, without losing generality, we shall impose the condition $|\varrho_n| < 1$, $\forall n > 1$.

The eigenfunctions of the Hamiltonian (10.5) are the customary functions (see Eq.(3.3))

$$H \chi = E \chi \longrightarrow \chi_\mp \propto \rho^{-1/2 \pm iE}. \quad (10.15)$$

From now on, we shall assume that E is a real number which is guaranteed by the self-adjointness of the Hamiltonian H . In the n^{th} interval we take

$$\chi_{\mp,n}(\rho) = e^{\pm i\pi/4} \frac{A_{\mp,n}}{\rho^{1/2 \mp iE}}, \quad \ell_n < \rho < \ell_{n+1}, \quad (10.16)$$




FIG. 10: Left: scattering process taking place at the mirror located at $\rho = \ell_n$ for $n > 1$ (Eq.(10.11)). Right: reflexion at the perfect mirror located at $\rho = \ell_1$ (Eq.(10.6)).

where $A_{\mp,n}$ are constants that, in general, will depend on E . The phases $e^{\pm i\pi/4}$ have been introduced by analogy with those appearing in Eq.(9.12). The boundary values of χ at $\rho = \ell_n^{\pm}$ ($n \geq 1$) are (see Eq.(10.7))

$$\chi_{\mp}(\ell_n^+) = \chi_{\mp,n}(\ell_n) = e^{\pm \frac{i\pi}{4}} \frac{A_{\mp,n}}{\ell_n^{1/2 \mp iE}}, \quad \chi_{\mp}(\ell_n^-) = \chi_{\mp,n-1}(\ell_n) = e^{\pm \frac{i\pi}{4}} \frac{A_{\mp,n-1}}{\ell_n^{1/2 \mp iE}}. \quad (10.17)$$

Let us define the vectors

$$|\mathbf{A}_n\rangle = \begin{pmatrix} A_{-,n} \\ A_{+,n} \end{pmatrix}, \quad n \geq 1. \quad (10.18)$$

The boundary conditions (10.6) together with Eqs.(10.17) imply

$$|\mathbf{A}_{n-1}\rangle = T_n |\mathbf{A}_n\rangle \quad (n \geq 2), \quad |\mathbf{A}_1\rangle = |\mathbf{A}_1(\vartheta)\rangle = \begin{pmatrix} 1 \\ e^{i\vartheta} \end{pmatrix}, \quad (10.19)$$

where the transfer matrix T_n is given by

$$T_n = \frac{1}{1 - |\varrho_n|^2} \begin{pmatrix} 1 + |\varrho_n|^2 & 2\varrho_n \ell_n^{-2iE} \\ 2\varrho_n^* \ell_n^{2iE} & 1 + |\varrho_n|^2 \end{pmatrix} \quad (n \geq 2). \quad (10.20)$$

The norm of the eigenstate can be computed using equations (10.4) and (10.16)

$$||\chi||^2 = \sum_{n=1}^{\infty} \log \frac{\ell_{n+1}}{\ell_n} \langle \mathbf{A}_n | \mathbf{A}_n \rangle, \quad \langle \mathbf{A}_n | \mathbf{A}_n \rangle = |A_{-,n}|^2 + |A_{+,n}|^2. \quad (10.21)$$

The log term comes from the integral of the norm of the wave function in the n^{th} interval, $\int_{\ell_n}^{\ell_{n+1}} d\rho/\rho$ (we used that E is real). If $\varrho_n = 0$ then $T_n = \mathbf{1}$ which implies that $|\mathbf{A}_{n-1}\rangle = |\mathbf{A}_n\rangle$. If this happens for all n , then $|\mathbf{A}_n\rangle = |\mathbf{A}_1\rangle$, in which case the norm of these states diverges, but they can be normalized using Dirac delta functions, so they correspond to scattering states. In the general case, iterating Eq.(10.19) yields $|\mathbf{A}_n\rangle$ in terms of $|\mathbf{A}_1(\vartheta)\rangle$

$$|\mathbf{A}_n\rangle = T_n^{-1} T_{n-1}^{-1} \cdots T_2^{-1} |\mathbf{A}_1(\vartheta)\rangle, \quad n \geq 2. \quad (10.22)$$

For special values of ℓ_n and ϱ_n one can find the exact expression of these amplitudes. An example is $\ell_n = e^{n/2}$, $\varrho_n = \text{cte}$ [55]. In order to make contact with the Riemann zeros, we shall consider a limit where the reflection coefficients vanish asymptotically.

Summary:

- ✓ The massless Dirac Hamiltonian with delta function potential is solvable by transfer matrix methods.
- ✓ The model is completely characterized by the set of parameters $\{\ell_n, \varrho_n\}_{n=2}^{\infty}$ and ϑ .

XI. HEURISTIC APPROACH TO THE SPECTRUM

Let us replace ϱ_n by $\varepsilon \varrho_n$, and consider the limit $\varepsilon \rightarrow 0$ of the transfer matrix (10.20)

$$T_n \simeq \mathbf{1} + \varepsilon \tau_n + O(\varepsilon^2), \quad \tau_n = \begin{pmatrix} 0 & 2\varrho_n \ell_n^{-2iE} \\ 2\varrho_n^* \ell_n^{2iE} & 0 \end{pmatrix} \quad (n \geq 2). \quad (11.1)$$




FIG. 11: Localization of the mirrors corresponding to the choice (11.4), together with the values of $\mu(n)$.




FIG. 12: Depiction of the amplitudes $A_{\pm, \infty}$ as the superposition of a principal wave with the waves resulting from the scattering with all the mirrors along its trajectory (see Eq.(11.3)). The terms of higher order in ε correspond to more than one scattering.

Plugging this equation into Eq.(10.22) yields

$$|\mathbf{A}_n\rangle \simeq \left(1 - \varepsilon \sum_{m=2}^n \tau_m\right) |\mathbf{A}_1(\vartheta)\rangle + O(\varepsilon^2), \quad n \geq 2, \quad (11.2)$$

and in components

$$A_{-,n} \simeq 1 - 2\varepsilon e^{i\vartheta} \sum_{m=2}^n \varrho_m \ell_m^{-2iE} + O(\varepsilon^2), \quad A_{+,n} \simeq e^{i\vartheta} - 2\varepsilon \sum_{m=2}^n \varrho_m^* \ell_m^{2iE} + O(\varepsilon^2). \quad (11.3)$$

For a normalizable state, the amplitudes $A_{\pm,n}$ have to vanish as $n \rightarrow \infty$. In the next section we shall study in detail the normalizability of the state. We shall make the following choice of lengths and reflection coefficients [55]

$$\ell_n = n^{1/2}, \quad \varrho_n = \frac{\mu(n)}{n^{1/2}}, \quad n > 1, \quad (11.4)$$

where $\mu(n)$ is the Moëbius function that is equal to $(-1)^r$, with r the number of distinct primes factors of a square free integer n , and $\mu(n) = 0$, if n is divisible by the square of a prime number [4]. See figs. 11 and 12 for a graphical representation of Eqs.(11.4) and (11.3). The Moëbius function has been used in the past to provide physical models of prime numbers, most notably in the ideal gas of primons with fermionic statistics [102]-[103] and a potential whose semiclassical spectrum are the primes [104]- [108].

Another motivation of the choice (11.4) is the following [55]. Consider a fermion that leaves the boundary at $\rho = \ell_1$, moves rightwards until it hits the mirror at $\rho = \ell_n$ where it gets reflected and returns to the boundary. The time lapse for the entire trajectory is given by

$$\tau_n = 2 \log(\ell_n / \ell_1) \quad (11.5)$$

where we used the Rindler metric Eq.(6.13). If the mirror is associated to the prime p , that is $\ell_p = \sqrt{p}$, the time will be given by $\tau_p = \log p$. This result reminds the Berry conjecture that postulates the existence of a classical chaotic Hamiltonian whose primitive periodic orbits are labelled by the primes p , with periods $\log p$, and whose quantization will give the Riemann zeros as energy levels [19]. A classical Hamiltonian with this property has not been found, but the array of mirrors presented above, displays some of its properties. In particular, the trajectory between the boundary and the mirror at ℓ_p , with p a prime number, behaves as a primitive orbit with a period $\log p$. Moreover, the trajectories and periods of these orbits are independent of the energy of the fermion because it moves at the speed of light.

Let us work out the consequences of the choice (11.4). The conditions for a normalizable eigenstate, that is $\lim_{n \rightarrow \infty} A_{\pm, n} = 0$, are

$$1 \simeq 2\varepsilon e^{i\vartheta} \sum_{n=1}^{\infty} \frac{\mu(n)}{n^{\frac{1}{2}+iE}} = \frac{2\varepsilon e^{i\vartheta}}{\zeta(\frac{1}{2}+iE)}, \quad 1 \simeq 2\varepsilon e^{-i\vartheta} \sum_{n=1}^{\infty} \frac{\mu(n)}{n^{\frac{1}{2}-iE}} = \frac{2\varepsilon e^{-i\vartheta}}{\zeta(\frac{1}{2}-iE)}, \quad (11.6)$$

where we have included the term $n = 1$ in the series because it does not modify its value when $\varepsilon \rightarrow 0$. We also employed the formula $\sum_{n=1}^{\infty} \mu(n)/n^s = 1/\zeta(s)$ for a value of s where the series may not converge. In the next section we shall compute the value of the finite sum that determines the norm of the state. Let $E_n(\varepsilon)$ be a solution of these equations and make the expansion in ε ,

$$E_n(\varepsilon) = E_n(0) + \varepsilon E'_n(0) + O(\varepsilon^2). \quad (11.7)$$

Plugging it into the zeta function gives

$$\zeta(\frac{1}{2} \pm iE_n(\varepsilon)) = \zeta(\frac{1}{2} \pm iE_n(0)) \pm i\varepsilon E'_n(0) \zeta'(\frac{1}{2} \pm iE_n) + O(\varepsilon^2), \quad (11.8)$$

where $\zeta'(s) = d\zeta(s)/ds$. The numerators of the fractions in Eq.(11.6) are of order ε , which forces the denominators to have the same order. In the limit $\varepsilon \rightarrow 0$ this is satisfied if

$$\zeta(\frac{1}{2} \pm iE_n) = 0, \quad E_n \equiv E_n(0), \quad (11.9)$$

and

$$1 = \frac{-2i e^{i\vartheta}}{E'_n(0) \zeta'(\frac{1}{2} + iE_n)}, \quad 1 = \frac{2i e^{-i\vartheta}}{E'_n(0) \zeta'(\frac{1}{2} - iE_n)}. \quad (11.10)$$

We have assumed that $\frac{1}{2} \pm iE_n$ is a simple zero of ζ , so $\zeta'(\frac{1}{2} \pm iE_n) \neq 0$. The last two equations lead to

$$\frac{\zeta'(\frac{1}{2} + iE_n)}{\zeta'(\frac{1}{2} - iE_n)} = -e^{2i\vartheta}, \quad E'_n(0) = \frac{-2i e^{i\vartheta}}{\zeta'(\frac{1}{2} + iE_n)}. \quad (11.11)$$

Using the Riemann-Siegel Z function, that appears in Eq.(2.9), one finds [109]

$$\zeta'(\frac{1}{2} \pm iE_n) = \mp i e^{\mp i\theta(E_n)} Z'(E_n), \quad (11.12)$$

where we have used Eq.(11.9). This equation allows one to write Eq.(11.11) as

$$e^{2i(\vartheta + \theta(E_n))} = 1, \quad (11.13)$$

and

$$E'_n(0) = \frac{2 e^{i(\vartheta + \theta(E_n))}}{Z'(E_n)}. \quad (11.14)$$

We can collect these results in the equation

$$\text{If } \zeta(\frac{1}{2} \pm iE_n) = 0 \text{ and } e^{2i(\vartheta + \theta(E_n))} = 1 \iff H_{\vartheta} \chi_{E_n} = E_n \chi_{E_n}. \quad (11.15)$$

Observe that ϑ is fixed mod π . In the next section we shall fix this ambiguity. This equation is heuristic. It has been derived by i) solving the eigenvalue equation in the limit $\varepsilon \rightarrow 0$, ii) imposing the vanishing of the eigenfunction




FIG. 13: Schematic representation of the array of mirrors that give rise to a spectral realization of the Riemann zeros. The red and blue lines represent the left and right wave functions $\chi_{\pm,n}(\rho)$. The wave functions are discontinuous at the moving mirrors located at the positions $\ell_n = \sqrt{n}$ with n a square free integer. The knob on the left represents the scattering phase at the perfect mirror that is set to minus the phase of the zeta function at the zero E_n , namely $\vartheta = -\theta(E_n) \bmod \pi$.

at infinity and iii) using the Dirichlet series of $1/\zeta(s)$ in a region where it may not converge. In the next section we shall derive Eq.(11.15) without making the previous assumptions (see Eq.(12.34)). Let us notice that this spectral realization of the *zeros* requires the fine tuning of the parameter of ϑ in terms of the phase of the zeta function, $\theta(E_n)$ (see fig. 13). This realization is different from the Pólya-Hilbert conjecture of a single Hamiltonian encompassing all the Riemann zeros at once. This Hamiltonian would exist if $\theta(E_n) = \theta_0, \forall n$, but this is certainly not the case.

Summary:

- ✓ A Riemann zero, on the critical line, becomes an eigenvalue of the Hamiltonian H_ϑ by tuning the phase ϑ according to the phase of the zeta function.
- ✗ The previous result is obtained in the limit $\varepsilon \rightarrow 0$ and is heuristic.

XII. THE RIEMANN ZEROS AS SPECTRUM AND THE RIEMANN HYPOTHESIS

In this section we provide more rigorous arguments that support the results obtained in the previous section. Let us first review the main properties of the model discussed so far. The Hamiltonian, Eq.(10.5), describes the dynamics of a massless Dirac fermion in the region of Rindler spacetime bounded by the hyperbola $\rho = \ell_1$. The reflection of the wave function at this boundary is characterized by a parameter ϑ , which is real for a self-adjoint Hamiltonian. At the positions $\ell_{n>1}$ the wave function is discontinuous due to the presence of delta functions potentials characterized by the reflections amplitudes ϱ_n , that provide the matching conditions of the wave function at those sites. An eigenfunction χ , with eigenvalue E , has a simple expression, Eq.(10.16), in terms of the amplitudes $A_{n,\pm}$, that are related by the transfer matrix T_n (10.20). The norm of χ is given by the sum of the squared length of the vectors \mathbf{A}_n , weighted with a factor that depends on the positions ℓ_n , Eq.(10.21). We introduce an scale factor ε in the parameters ϱ_n , that allows us to study the limit $\varepsilon \rightarrow 0$, where the mirrors become transparent. In this way we found an ansatz for the parameters ℓ_n and ϱ_n that heuristically led to an individual spectral realization of the *zeros* by fine tuning the parameter ϑ .

1. Normalizable eigenstates

Under the choice $\ell_n = n^{1/2}$, Eq.(10.21) becomes

$$\|\chi\|^2 = \frac{1}{2} \sum_{n=1}^{\infty} \log \left(1 + \frac{1}{n} \right) \langle \mathbf{A}_n | \mathbf{A}_n \rangle. \quad (12.1)$$

This series can be replaced by

$$\|\chi\|_c^2 \equiv \sum_{n=1}^{\infty} \frac{1}{n} \langle \mathbf{A}_n | \mathbf{A}_n \rangle, \quad (12.2)$$

which is convergent if and only if (12.1) is convergent. The vectors \mathbf{A}_n are obtained by acting on $\mathbf{A}_1(\vartheta)$ with the transfer matrices T_n (see Eq.(10.22)). These matrices have unit determinant and can be written as the exponential of traceless hermitean matrices, that is,

$$T_n = e^{\tau_n}, \quad \tau_n = \begin{pmatrix} 0 & r_n \ell_n^{-2iE} \\ r_n^* \ell_n^{2iE} & 0 \end{pmatrix}, \quad \forall E \in \mathbb{R}, \quad (12.3)$$

where taking $|\varrho_n| < 1$,

$$r_n = \frac{\varrho_n}{|\varrho_n|} \log \frac{1 + |\varrho_n|}{1 - |\varrho_n|}, \quad \varrho_n = \frac{r_n}{|r_n|} \tanh \frac{|r_n|}{2}. \quad (12.4)$$

To derive Eq.(12.3) we have used the relation

$$\exp \begin{pmatrix} 0 & a \\ b & 0 \end{pmatrix} = \begin{pmatrix} \cosh(\sqrt{ab}) & \frac{a}{\sqrt{ab}} \sinh(\sqrt{ab}) \\ \frac{b}{\sqrt{ab}} \sinh(\sqrt{ab}) & \cosh(\sqrt{ab}) \end{pmatrix}, \quad \forall a, b \in \mathbb{C} - \{0\}. \quad (12.5)$$

If $|\varrho_n| \ll 1$ one gets $r_n \simeq 2\varrho_n$, hence in that limit both parameters give the same result. Using Eq.(12.3), the recursion relation (10.22) reads

$$|\mathbf{A}_k\rangle = e^{-\tau_k} e^{-\tau_{k-1}} \dots e^{-\tau_2} |\mathbf{A}_1\rangle, \quad k \geq 2. \quad (12.6)$$

2. The Magnus expansion

In general, the product of matrices in Eq.(12.6) is very difficult to compute. However, we can make the approximation introduced in the previous section were the coefficients r_n are replaced by εr_n , with $\varepsilon > 0$, and then take the limit $\varepsilon \rightarrow 0$. Under this replacement Eq.(12.6) becomes

$$|\mathbf{A}_k\rangle = e^{-\varepsilon\tau_k} e^{-\varepsilon\tau_{k-1}} \dots e^{-\varepsilon\tau_2} |\mathbf{A}_1\rangle \quad (k \geq 2). \quad (12.7)$$

The product of exponentials of matrices can be expressed as the exponential of a matrix given by the Magnus expansion [110]

$$e^{-\varepsilon\tau_k} e^{-\varepsilon\tau_{k-1}} \dots e^{-\varepsilon\tau_2} = \exp \left(-\varepsilon \sum_{n=2}^k \tau_n - \frac{\varepsilon^2}{2} \sum_{n_1 > n_2 = 2}^k [\tau_{n_1}, \tau_{n_2}] + O(\varepsilon^3) \right) \quad (k \geq 2). \quad (12.8)$$

In the limit $\varepsilon \rightarrow 0$ we shall make the approximation

$$e^{-\varepsilon\tau_k} e^{-\varepsilon\tau_{k-1}} \dots e^{-\varepsilon\tau_2} \simeq \exp \begin{pmatrix} 0 & -\varepsilon \sum_{n=2}^k r_n \ell_n^{-2iE} \\ -\varepsilon \sum_{n=2}^k r_n^* \ell_n^{2iE} & 0 \end{pmatrix} \simeq \exp \begin{pmatrix} 0 & -\varepsilon M_z(k) \\ -\varepsilon M_z^*(k) & 0 \end{pmatrix}, \quad (12.9)$$

that with the choice (recall (11.4))

$$r_n = \frac{\mu(n)}{n^{1/2}} \quad (12.10)$$

gives

$$M_z(k) = 1 + \sum_{n=2}^k r_n \ell_n^{-2iE} = \sum_{n=1}^k \frac{\mu(n)}{n^z}, \quad z = \frac{1}{2} + iE. \quad (12.11)$$

We have added the constant 1 to $M_z(k)$, that does not affect the results in the limit $\varepsilon \rightarrow 0$. Using Eqs.(12.5) and (12.9) into (12.7) yields

$$|\mathbf{A}_n\rangle \simeq e^{i(\vartheta - \Phi_z(n))} \begin{pmatrix} e^{-|\varepsilon M_z(n)|} \cos(\frac{1}{2}(\vartheta - \Phi_z(n))) - i e^{|\varepsilon M_z(n)|} \sin(\frac{1}{2}(\vartheta - \Phi_z(n))) \\ e^{-|\varepsilon M_z(n)|} \cos(\frac{1}{2}(\vartheta - \Phi_z(n))) + i e^{|\varepsilon M_z(n)|} \sin(\frac{1}{2}(\vartheta - \Phi_z(n))) \end{pmatrix} \quad (n \geq 2), \quad (12.12)$$

whose norm (12.2) is

$$\|\chi\|_c^2 \simeq \sum_{n=1}^{\infty} \frac{1}{n} \left[e^{-2|\varepsilon M_z(n)|} (1 + \cos(\vartheta - \Phi_z(n))) + e^{2|\varepsilon M_z(n)|} (1 - \cos(\vartheta - \Phi_z(n))) \right] \quad (12.13)$$

where

$$e^{-i\Phi_z(n)} = \frac{M_z(n)}{|M_z(n)|} \quad (12.14)$$

3. The Mertens function and the Perron formula

Setting $z = 0$ in Eq.(12.11) gives the Mertens function

$$M(x) = M_0(x) = \sum_{1 \leq n \leq x} \mu(n). \quad (12.15)$$

where the argument has been extended to include non integer values. This function has played an important role in Number Theory. In 1897 Mertens conjectured that $|M(x)| \leq x^{1/2}$, $\forall x > 1$ [111]. If this conjecture were true it would imply the RH. However in 1985, Odlyzko and the Riele proved that this conjecture is false [112]. In fact, the RH is equivalent to a weaker statement, namely [3]

$$M(x) = O(x^{1/2+c}), \quad \forall c > 0. \quad (12.16)$$

But, even if the RH is true, the rate of growth of $M(x)$ is unknown [112]. Gonek in 1990 and Ng in 2004 have conjectured that $M(x)$ satisfies [113]

$$M(x) = O(x^{1/2}(\log \log \log x)^{5/4}), \quad (12.17)$$

using the big O notation [114]. In the rest of this section we shall investigate the conditions under which a *zero* of the zeta function is an eigenvalue of the Hamiltonian introduced in the section X, and what is the relation of this problem with the RH.

To compute $M_z(n)$ it is convenient to define the function

$$M'_z(x) \equiv \sum'_{1 \leq n \leq x} \frac{\mu(n)}{n^z}, \quad (12.18)$$

where $\sum'_{1 \leq n \leq x}$ means that the last term in the sum is multiplied by 1/2 when x is an integer. Fig. 14 shows $M'_z(n)$ as a function of E for several values of n . Observe that $|M'_z(n)|$ increases with n when E is a *zero*. We shall derive below this behavior. To compute $M'_z(x)$ we use the Perron's formula [114, 115]

$$M'_z(x) = \lim_{T \rightarrow \infty} \int_{c-iT}^{c+iT} \frac{ds}{2\pi i} \frac{1}{\zeta(s+z)} \frac{x^s}{s}, \quad c > \frac{1}{2}, \quad (12.19)$$

where we have used that $\text{Re } z = 1/2$. The integral (12.19) can be done by residue calculus [55]

$$\lim_{T \rightarrow \infty} \int_{c-iT}^{c+iT} \frac{ds}{2\pi i} F(s) = \sum_{\text{Re } s_j < c} \text{Res}_{s_j} F(s), \quad F(s) = \frac{1}{\zeta(s+z)} \frac{x^s}{s}, \quad (12.20)$$

where the sum runs over the poles s_j of $F(s)$ located to the left of the line of integration $\text{Re } s = c$, that is $\text{Re } s_j < c$. The poles of $F(s)$ come from the zeros of $s\zeta(s+z)$. The pole at $s = 0$ can be simple, or multiple, depending on the values of $\zeta(z)$ and its derivatives. The other poles of $F(s)$ come from the zeros of $\zeta(s+z)$, say $s_j + z = \rho_j$, and they all contribute to the residue of $F(s)$ because they lie to the left of the integration line,

$$\text{Re } s_j = \text{Re}(\rho_j - z) = \text{Re } \rho_j - \frac{1}{2} < \frac{1}{2} < c. \quad (12.21)$$

Recall that all the *zeros* of ζ , trivial and non trivial, satisfy $\text{Re } \rho_j < 1$. To compute the residues of Eq.(12.20) we consider three possible cases: $s = 0$, $s_j + z$ a trivial zero of ζ and $s_j + z$ a non trivial zero of ζ :




FIG. 14: Plot of $|M'_z(n)|$ computed doing the sum in Eq.(12.18), for $E \in (10, 23)$ and $n = 50, 100, 150$ (blue, orange, red curves) and $1/|\zeta(1/2 + iE)|$ (dotted line). Observe the increase with n at $E = 14.13\dots$ and $E = 21.02\dots$ which are the first two *zeros* of ζ .

- $s = 0$. Let $m \geq 0$ be the lowest integer such that $\zeta^{(m)}(z) = d^m \zeta(z)/dz^m \neq 0$. Then $F(s)$ has a pole of order $m + 1$ at $s = 0$ with residue [116]

$$\text{Res}_{s=0} F(s) = \begin{cases} 1/\zeta(z) & \text{if } \zeta(z) \neq 0, \\ \log x / \zeta'(z) - \frac{1}{2} \zeta''(z) / (\zeta'(z))^2 & \text{if } \zeta(z) = 0, \zeta'(z) \neq 0. \\ \vdots & \vdots \\ (\log x)^m / \zeta^{(m)}(z) + O((\log x)^{m-1}) & \text{if } \zeta(z) = 0, \dots, \zeta^{(m-1)}(z) = 0, \zeta^{(m)}(z) \neq 0. \end{cases} \quad (12.22)$$

- $s_n = -2n - z$ ($n = 1, 2, \dots$), where $F(s)$ has a simple pole due to the trivial zeros of $\zeta(s + z)$.

$$\text{Res}_{s=-2n-z} F(s) = \frac{x^{-2n-z}}{-(2n+z)\zeta'(-2n)}, \quad n = 1, 2, \dots, \infty. \quad (12.23)$$

- $s_j = \rho_j - z \neq 0$, where $F(s)$ has a pole due to the non trivial zero ρ_j of ζ that we shall assume is simple,

$$\text{Res}_{s=\rho_j-z} F(s) = \frac{x^{\rho_j-z}}{(\rho_j - z)\zeta'(\rho_j)}, \quad 0 < \text{Re } \rho_j < 1, \quad \rho_j \neq z. \quad (12.24)$$

If ρ_j is a double *zero* then we get $2(\log x)x^{\rho_j-z}/(\rho_j - z)\zeta''(z)$ plus corrections. To simplify the discussion we shall suppose that all the *zeros* are simple.

Collecting these results we have

$$M_z(x) = \frac{1}{\zeta(z)} + \sum_{\rho_j} \frac{x^{\rho_j-z}}{(\rho_j - z)\zeta'(\rho_j)} + \sum_{n=1}^{\infty} \frac{x^{-2n-z}}{-(2n+z)\zeta'(-2n)}, \quad \text{if } \zeta(z) \neq 0, \quad (12.25)$$

$$M_z(x) = \frac{\log x}{\zeta'(z)} - \frac{\zeta''(z)}{2(\zeta'(z))^2} + \sum_{\rho_j \neq z} \frac{x^{\rho_j-z}}{(\rho_j - z)\zeta'(\rho_j)} + \sum_{n=1}^{\infty} \frac{x^{-2n-z}}{-(2n+z)\zeta'(-2n)}, \quad \text{if } \zeta(z) = 0, \zeta'(z) \neq 0. \quad (12.26)$$

where the sum \sum_{ρ_j} runs over the non trivial zeros of ζ . These equations are verified numerically in Fig.15. The Mertens function $M'_0(x) = \sum'_{1 \leq n \leq x} \mu(n)$, can be computed similarly obtaining the expression [3, 115]

$$M_0(x) = -2 + \sum_{\rho_j} \frac{x^{\rho_j}}{\rho_j \zeta'(\rho_j)} + \sum_{n=1}^{\infty} \frac{x^{-2n}}{-2n \zeta'(-2n)}, \quad (12.27)$$

that, using the fact that $\zeta(1/2) = -1/2$, coincides with Eq.(12.25) for $z = 0$. The term in these equations associated to the trivial *zeros* converges quickly and is finite for all x due to the exponential increase of $\zeta'(-2n)$ [10]

$$\zeta'(-2n) = \frac{(-1)^n \zeta(2n+1)(2n)!}{2^{2n+1} \pi^{2n}} \xrightarrow{n \rightarrow \infty} (-1)^n \sqrt{\pi n} \left(\frac{n}{e\pi} \right)^{2n}. \quad (12.28)$$




FIG. 15: Plot of $|M'_z(n)|$ for $n = 10, \dots, 50$ and $E = 20$ (left) and $E = 14.13..$ (right). In red the values obtained doing the sum in Eq.(12.18). In blue the sum of Eq.(12.25) for $E = 20$ and Eq.(12.26) for $E = 14.13..$, including the first 100 Riemann zeros, and 20 trivial zeros. Observe the accuracy of the approximation. The slow increase in the latter plot is due to the factor $\log n$ in Eq.(12.26).

A comparison of Eqs.(12.25) and (12.26) with Eq.(12.27) suggests the following relations between the order of the functions $M_z(x)$ and $M(x)$, as a function of $z = 1/2 + iE, E \in \mathbb{R}$

$$\begin{aligned} M_z(x) &= O(x^{-1/2}M(x)), & \text{if } \zeta(z) \neq 0 \\ M_z(x) &= O(\log x + x^{-1/2}|M(x)|), & \text{if } \zeta(z) = 0, \zeta'(z) \neq 0. \end{aligned} \quad (12.29)$$

Let us consider the two possibilities concerning the truth of the RH:

- If the RH is true then equations (12.16) and (12.29) imply that

$$\begin{aligned} M_z(x) &= O(x^c), & \text{if } \zeta(z) \neq 0, \\ M_z(x) &= O(x^c), & \text{if } \zeta(z) = 0, \zeta'(z) \neq 0. \end{aligned} \quad (12.30)$$

This information is insufficient to determine the convergence of the norm (12.13) and does not distinguish between *zeros* and non *zeros*. Assuming the Gonek-Ng conjecture (12.17) one finds that

$$\begin{aligned} M_z(x) &= O((\log \log \log x)^{5/4}), & \text{if } \zeta(z) \neq 0, \\ M_z(x) &= O(\log x), & \text{if } \zeta(z) = 0, \zeta'(z) \neq 0. \end{aligned} \quad (12.31)$$

The latter equation, together with (12.26), imply

$$M_x(z) \rightarrow \frac{\log x}{\zeta'(z)} = \frac{i e^{i\theta(E)} \log x}{Z'(E)} \quad \text{as } x \rightarrow \infty, \quad \text{if } \zeta(z) = 0 \text{ and } \zeta'(z) \neq 0, \quad (12.32)$$

hence $\Phi_z(n)$, given in Eq.(12.14), behaves as

$$e^{-i\Phi_z(n)} \rightarrow i e^{i\theta(E)} \text{sign } Z'(E) \quad \text{as } n \rightarrow \infty, \quad (12.33)$$

whose limit is independent on n . Thus we can choose ϑ in such a way that the term proportional to $e^{2\varepsilon|M_z(n)|}$ in Eq.(12.13) vanishes in the limit $n \rightarrow \infty$, namely

$$\vartheta = \lim_{n \rightarrow \infty} \Phi_z(n) = - \left(\theta(E) + \frac{\pi}{2} \text{sign } Z'(E) \right). \quad (12.34)$$

Notice that ϑ depends on $\theta(E)$ and the sign of $Z'(E)$, a feature that is left undetermined in Eq.(11.13). The norm (12.13) then becomes

$$\|\chi\|_c^2 \simeq \sum_{n=1}^{\infty} \frac{2}{n} e^{-2\varepsilon \log n / |Z'(E)|} = 2\zeta \left(1 + \frac{2\varepsilon}{|Z'(E)|} \right) < \infty, \quad (12.35)$$

that is finite for all $\varepsilon > 0$. This result shows that a *zero* of the zeta function gives a normalizable state, in agreement with heuristic derivation proposed in the previous section, but there are some differences. First of

all, the eigenvalue E does not need to be expanded in series of ε (recall Eq.(11.7)). It is taken to be a *zero* of ζ from the beginning. This choice generates the $\log x$ term in Eq.(12.26) and is responsible for the finiteness of the norm after the appropriate choice of the phase (12.34), that also differs from the heuristic value (11.13). On the other hand, if ϑ does not satisfy Eq.(12.34), then the norm of the state will diverge badly and so the *zero* E will be missing in the spectrum. Finally, if E is not a *zero*, we expect that the state will belong generically to the continuum. Fig. 16 shows the expected spectrum of the model, which recalls the Connes scenario of missing spectral lines, except that in our case, one can pick up a zero at a time by tuning ϑ .




FIG. 16: Graphical representation of the spectrum of the model. It is expected to consists of an infinite number of bands separated by forbidden regions of width proportional to ε . The latter regions may contain a *zero* E_n if the phase ϑ is chosen according to Eq.(12.34). Otherwise the *zeros* will be missing in the spectrum, as illustrated by the points E_{n-1} and E_{n+1} .

- If the RH is false then there will be at least four *zeros* outside the critical line, say $\rho_c = \sigma_c + iE_c, \bar{\rho}_c = \sigma_c - iE_c, 1 - \rho_c$ and $1 - \bar{\rho}_c$, with $\sigma_c > \frac{1}{2}, E_c \in \mathbb{R}_+$. We shall choose the highest value of σ_c . The asymptotic behavior of $M_z(x)$ will be given by the *zeros* located to the right of the critical line,

$$M_z(x) \rightarrow \frac{x^{\rho_c - z}}{(\rho_c - z)\zeta'(\rho_c)} + \frac{x^{\bar{\rho}_c - z}}{(\bar{\rho}_c - z)\zeta'(\bar{\rho}_c)} \quad \text{as } x \rightarrow \infty. \quad (12.36)$$

To simplify the discussion let us choose $E \gg E_c$, which yields the approximation

$$M_z(x) \rightarrow \frac{2i x^{\sigma_c - 1/2 - iE}}{E|\zeta'(\rho_c)|} \cos(E \log x - \phi_c) \quad \text{as } x \rightarrow \infty, \quad (12.37)$$

where $e^{i\phi_c} = \zeta'(\rho_c)/|\zeta'(\rho_c)|$. The phase $\Phi_z(n)$ is given by Eq.(12.14)

$$\Phi_z(n) \rightarrow E \log n - \frac{\pi}{2} \text{sign}(\cos(E \log n - \phi_c)) \quad \text{as } n \rightarrow \infty. \quad (12.38)$$

Correspondingly, the norm (12.13) diverges so badly, $\propto \sum_n \frac{1}{n} \exp(Cn^{\sigma_c - 1/2}) \dots$, for any value of ϑ , that the state will not be normalizable even using Dirac delta functions. This result occurs for all eigenenergies E . Therefore the Hamiltonian will not admit a spectral decomposition, but this is impossible because it is a well defined self-adjoint operator. We conclude that a *zero* outside the critical line does not exist which provides an argument likely to be persuasive to physicists for the truth of the Riemann hypothesis.

XIII. THE RIEMANN INTERFEROMETER

The model considered in the previous sections looks at first glance quite difficult to simulate. We shall next show that this model is equivalent to another one that can be implemented in the Lab. We shall call this system the Riemann interferometer. The basic idea can be illustrated with the mapping between the quantum xp hamiltonian and the momentum operator p . Let us make the change of coordinates $x = \log \rho$ and relate the wave functions in both coordinates, $\phi(x)$ and $\psi(\rho)$, as follows

$$\phi(x) = \left(\frac{d\rho}{dx} \right)^{1/2} \psi(\rho) = e^{x/2} \psi(e^x). \quad (13.1)$$

An eigenstate of the Hamiltonian $(\rho \hat{p}_\rho + \hat{p}_\rho \rho)/2$, with eigenvalue E , is mapped by Eq.(13.1) into an eigenstate of the momentum operator $\hat{p}_x = -i\partial_x$ with the same eigenvalue,

$$\psi(\rho) = \frac{1}{\rho^{1/2 - iE}} \implies \phi(x) = e^{iEx}. \quad (13.2)$$

This shows that the energy E can also be seen as momentum. For a relativistic massless fermion, this is always the case. The measure that defines the scalar product of the corresponding Hilbert spaces are one-to-one related

$$\int_{\ell}^{\infty} d\rho \psi_1^*(\rho) \psi_2(\rho) = \int_{\log \ell}^{\infty} dx \phi_1^*(x) \phi_2(x). \quad (13.3)$$

The operator $(\rho \hat{p}_{\rho} + \hat{p}_{\rho} \rho)/2$ is self-adjoint in the interval $(0, \infty)$ but not in the interval $(1, \infty)$, just like \hat{p}_x is self-adjoint in the real line $(-\infty, \infty)$ but not in the half line $(0, \infty)$ [35, 97]. The former case corresponds to the value $\ell = 0$ and the latter one to $\ell = 1$ in Eq.(13.3). Let us now consider the Dirac Hamiltonian in the Rindler variable ρ , given in Eq.(10.5). It becomes in the x variable

$$H = \begin{pmatrix} -i\partial_x & 0 \\ 0 & i\partial_x \end{pmatrix}. \quad (13.4)$$

Unlike \hat{p}_x , this Hamiltonian is self-adjoint in the interval $x \in (\log \ell_1, \infty)$. We choose for convenience $\ell_1 = 1$. The moving mirrors located at $\rho = \ell_n$ are now placed at the positions $x = d_n$, with $d_n = \log \ell_n$, so for $\ell_n = \sqrt{n}$, we get

$$d_n = \frac{1}{2} \log n, \quad (13.5)$$

where n are square free integers and the reflection coefficients are given by $r_n = \mu(n)/\sqrt{n}$. Fig. 17 shows the array of mirrors satisfying Eq.(13.5). One can easily generalize this interferometer to provide a spectral realization of the *zeros* of Dirichlet L -functions, by changing the reflection coefficients r_n ,

$$L_{\chi}(s) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s} \longrightarrow r_n = \frac{\mu(n) \chi(n)}{n^{1/2}}, \quad (13.6)$$

where $\chi(n)$ is the Dirichlet character associated to the L -function. It would be interesting to replace the massless fermions by massless bosons, say photons and study what kind of Riemann interferometer arise.




FIG. 17: Graphical representation of the array of mirrors in Minkowski space that reproduce the Riemann zeros. The phase at the boundary ϑ has to be chosen according to Eq.(12.34) in order that E is an eigenvalue of the Hamiltonian. Recall Fig. 13. Between the mirrors the wave functions are plane waves.

XIV. CONCLUSIONS

In this paper we have reviewed the spectral approach to the RH that started with the Berry-Keating-Connes xp model and continued with several works aimed to provide a physical realization of the Riemann zeros. The main steps in this approach are: i) spectral realization of Connes's xp model using the Landau model of an electron in a magnetic field and electrostatic potential, ii) construction of modified quantum xp models whose spectra reproduce, in average, the behavior of the *zeros*, iii) reformulation of the $x(p + 1/p)$ model as a relativistic theory of a massive Dirac fermion in a region of Rindler space-time, iv) inclusion of the prime numbers into the massless Dirac equation by means of delta function potentials acting as moving mirrors that, in the limit where they become transparent, leads to a spectral realization of the *zeros*, v) a proof of the Riemann Hypothesis, and vi) proposal of an interferometer that may provide an experimental observation of the zeros of the Riemann zeta function and other Dirichlet L -functions.

The Pólya-Hilbert (PH) conjecture was proposed as a physical explanation of the RH based on the spectral properties of self-adjoint operators: there exists a *single* quantum Hamiltonian containing *all* the Riemann zeros in its spectrum which are therefore real numbers. This statement can be called the *global* version of the PH conjecture. Instead of this, we have found a *local* version according to which a Riemann zero E_n becomes an eigenvalue of the Hamiltonian H_ϑ provided the parameter ϑ , that characterizes the self-adjoint extension, is fine tuned to the combination $\theta(E_n) + \frac{\pi}{2}\text{sign}Z'(E_n)$. In this sense the Hamiltonian provides a physical realization of $\zeta(\frac{1}{2} + it)$, and not only of the Riemann-Siegel Z function. We have also obtained a proof of the RH by contradiction: the existence of a *zero* off the critical line implies that the eigenstates of H_ϑ are non normalizable in the discrete or continuum sense, which is impossible since H_ϑ is a self-adjoint operator. These results are obtained in the limit where the mirrors become transparent and assumes the convergence of some mathematical series that need to be analyzed more thoroughly. Finally, we have proposed an interferometer made of fermions propagating in a array of mirrors that may yield an experimental observation of the Riemann zeros in the Lab.

Acknowledgements

I am grateful for fruitful discussions and comments to Manuel Asorey, Michael Berry, Ignacio Cirac, Charles Creffield, Jon Keating, José Ignacio Latorre, André LeClair, Miguel Angel Martín-Delgado, Javier Molina-Vilaplana, Javier Rodríguez-Laguna, Mark Srednicki and Paul Townsend. I thanks Denis Bernard for pointing out an error in the first version of this manuscript. This work has been supported by the MINECO grant FIS2012-33642, Comunidad de Madrid grant QUITEMAD+ S2013/ICE-2801 and the grant SEV-2012-0249 of the "Centro de Excelencia Severo Ochoa" Programme.

-
- [1] B. Riemann, "On the number of primes less than a given magnitude", translated by H. M. Edwards from "Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse" (1859) in reference [2].
 - [2] H.M. Edwards, *Riemann's Zeta Function*, Academic Press, New York, 1974.
 - [3] E.C. Titchmarsh, *The Theory of the Riemann Zeta Function*, Oxford Univ. Press, Oxford, 1986.
 - [4] H. Davenport, *Multiplicative Number Theory*, Grad. Texts in Math., vol. **74**, Springer-Verlag, New York, 2000.
 - [5] E. Bombieri, Problems of the Millennium: The Riemann Hypothesis, Clay Mathematics Institute, (2000).
 - [6] P. Sarnak, Problems of the Millennium: The Riemann Hypothesis, Clay Mathematics Institute, (2004).
 - [7] B. Conrey, "The Riemann Hypothesis, Notices of AMS, March, 341 (2003).
 - [8] A. Ivić, "On some reasons for doubting the Riemann hypothesis", arXiv:math/0311162.
 - [9] G. J. Chaitin, "Thoughts on the Riemann hypothesis", arXiv:math/0306042.
 - [10] P. Borwein, S. Choi, B. Rooney and A. Weirathmueller (Eds.). *The Riemann Hypothesis. A Resource for the Afficionado and Virtuoso Alike*, CMS Books in Mathematics, Springer, Canada, 2008.
 - [11] A. Connes, "An essay on the Riemann Hypothesis", arXiv:1509.05576.
 - [12] G. Pólya, unpublished (c. 1914). See A. Odlyzko, Correspondence about the origins of the Hilbert - Pólya conjecture, <http://www.dtc.umn.edu/~odlyzko/polya/index.html> (1981-1982).
 - [13] H.L. Montgomery, "The pair correlation of the zeta function", Proc. Symp. Pure Math. **24**,181 (1973).
 - [14] A.M. Odlyzko, "Supercomputers and the Riemann zeta function", in Supercomputing 89: Supercomputing Structures & Computations, Proc. 4-th Intern. Conf. on Supercomputing, International Supercomputing Institute, St. Petersburg, FL, 348 (1989).
 - [15] A. M. Odlyzko, The 1020-th zero of the Riemann Zeta Function and 70 Million of its Neighbors, www.dtc.umn.edu/~odlyzko/unpublished, Preprint (1989).
 - [16] O. Bohigas, M. J. Gianonni and C. Schmit, "Characterization of chaotic quantum spectra and universality of level fluctuation". Phys. Rev. Lett. **52**, 1 (1984).
 - [17] M. V. Berry, "Semiclassical formula for the number variance of the Riemann zeros". Nonlinearity **1**, 399 (1988).
 - [18] M. L. Mehta, *Random Matrices*, Elsevier Ltd. The Netherlands, 2004.
 - [19] M. V. Berry, "Riemann's zeta function: a model for quantum chaos?", in Quantum Chaos and Statistical Nuclear Physics, edited by T. H. Seligman and H. Nishioka, Springer Lecture Notes in Physics Vol. **263**, p. 1, Springer, New York (1986).
 - [20] M. V. Berry, "Quantum chaology" (The Bakerian lecture), R. Soc. London, Ser. A **413**, 183 (1987).
 - [21] M. C. Gutzwiller, *Chaos in Classical and Quantum Mechanics*, SpringerVerlag, New York, 1990.
Bogomolny, E.B. and Keating, J.P.: Random matrix theory and the Riemann zeros I; three- and four-point correlations. Nonlinearity **8**, 11151131 (1995)

- [22] J. P. Keating, "Periodic orbits, spectral statistics and the Riemann zeros", in *Supersymmetry and Trace Formulae: Chaos and Disorder*, eds. I.V. Lerner, J.P. Keating & D.E Khmelnitskii (Plenum Press), 1-15 (1999).
- [23] J.P. Keating and N.C. Snaith, "Random matrix theory and $\zeta(1/2 + it)$ ", *Commun. Math. Phys.* **214**, 57 (2000).
- [24] P. Leboeuf, A. G. Monastera, O. Bohigas, "The Riemannium", *Reg. Chaot. Dyn.* **6**, 205 (2001).
- [25] E. Bogomolny, "Quantum and Arithmetical Chaos", in *Frontiers in Number Theory, Physics and Geometry*, Les Houches, 2003; arXiv:nlin/0312061.
- [26] Y. V. Fyodorov, G. A. Hiary, J. P. Keating, "Freezing Transition, Characteristic Polynomials of Random Matrices, and the Riemann Zeta-Function", *Phys. Rev. Lett.* **108**, 170601 (2012); arXiv:1202.4713.
- [27] L. A. Bunimovich and C. P. Dettmann, "Open circular billiards and the Riemann hypothesis", *Phys. Rev. Lett.* **94**, 100201 (2005); arXiv:nlin/0408009.
- [28] A. Selberg, "Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series", *Journal of the Indian Mathematical Society* **20**, 47 (1956).
- [29] D. Hejhal, "The Selberg trace formula and the Riemann zeta function". *Duke Math. J.* **43**, 441 (1976).
- [30] M.V. Berry, J.P. Keating, " $H = xp$ and the Riemann zeros", in *Supersymmetry and Trace Formulae: Chaos and Disorder*, ed. J.P. Keating, D.E. Khmelnitskii and I. V. Lerner, Kluwer, 1999.
- [31] M. V. Berry, J. P. Keating, "The Riemann zeros and eigenvalue asymptotics", *SIAM Review* **41**, 236, 1999.
- [32] A. Connes, "Trace formula in noncommutative geometry and the zeros of the Riemann zeta function", *Selecta Mathematica New Series* **5** 29, (1999); math.NT/9811068.
- [33] B. Aneva, "Symmetry of the Riemann operator", *Phys. Lett. B* **450**, 388 (1999).
- [34] G. Sierra, "The Riemann zeros and the cyclic renormalization group", *J. Stat. Mech.: Theor. Exp.* (2005) P12006; math.NT/0510572.
- [35] G. Sierra, " $H = xp$ with interaction and the Riemann zeros", *Nucl. Phys. B* **776**, 327 (2007); math-ph/0702034.
- [36] J. Twamley and G. J. Milburn, "The quantum Mellin transform", *New J. Phys.* **8**, 328 (2006); quant-ph/0702107.
- [37] G. Sierra, "Quantum reconstruction of the Riemann zeta function", *J. Phys. A: Math. Theor.* **40** (2007) 1; math-ph/0711.1063.
- [38] G. Sierra, "A quantum mechanical model of the Riemann zeros", *New J. Phys.* **10**, 033016 (2008); arXiv:0712.0705.
- [39] J. C. Lagarias, "The Schrödinger operator with Morse potential on the right half line", *Communications in Number Theory and Physics* **3** (2009), 323; arXiv:0712.3238.
- [40] J-F. Burnol, "On some bound and scattering states associated with the cosine kernel", arXiv:0801.0530.
- [41] G. Sierra and P.K. Townsend, "The Landau model and the Riemann zeros", *Phys. Rev. Lett.* **101**, 110201 (2008); arXiv:0805.4079.
- [42] S. Endres and F. Steiner, "The Berry-Keating operator on $L^2(R_+, dx)$ and on compact quantum graphs with general self-adjoint realizations", *J. Phys. A: Math. Theor.* **43**, 095204 (2010); arXiv:0912.3183.
- [43] G. Regniers and J. Van der Jeugt, "The Hamiltonian $H = xp$ and classification of $osp(1|2)$ representations", *Workshop Lie Theory and Its Applications in Physics VIII* (Varna, 2009); arXiv:1001.1285.
- [44] G. Sierra, "A Physics pathway to the Riemann hypothesis", Eds M. Asorey, et al. J. Vicente García Esteve, M. F. Rañada, J. Sesma, *Prensas Univ. Zaragoza*, Spain (2009); arXiv:1012.4264.
- [45] G. Sierra and J. Rodríguez-Laguna, "The $H = xp$ model revisited and the Riemann zeros". *Phys. Rev. Lett.* **106**, 200201 (2011); arXiv:1102.5356.
- [46] M. Srednicki, "The Berry-Keating Hamiltonian and the Local Riemann Hypothesis", *J. Phys. A: Math. Theor.* **44** 305202 (2011); arXiv:1104.1850.
- [47] M. Srednicki, "Nonclassical Degrees of Freedom in the Riemann Hamiltonian", *Phys. Rev. Lett.* **107**, 100201 (2011); arXiv:1105.2342.
- [48] G. Sierra, "General covariant xp models and the Riemann zeros", *J. Phys. A: Math. Theor.* **45** 055209 (2012); arXiv:1110.3203.
- [49] M. V. Berry and J. P. Keating, "A compact hamiltonian with the same asymptotic mean spectral density as the Riemann zeros", *J. Phys. A: Math. Theor.* **44**, 285203 (2011).
- [50] K. S. Gupta, E. Harikumar and A. R. de Queiroz, "A Dirac type xp -Model and the Riemann Zeros", *Eur. Phys. Lett.* **102** 10006 (2013); arXiv:1205.6755.
- [51] J. Molina-Vilaplana and G. Sierra, "An xp model on AdS_2 spacetime", *Nucl. Phys. B* **877**, 107 (2013); arXiv:1212.2436.
- [52] M. C. Nucci, "Spectral realization of the Riemann zeros by quantizing $H = w(x)(p + \ell_p^2/p)$: the Lie-Noether symmetry approach", *Journal of Physics: Conference Series* **482**, 012032 (2014).
- [53] J. C. Andrade, "Hilbert-Pólya conjecture, zeta-functions and bosonic quantum field theories", *Int. J. Mod.Phys. A* **28**, 1350072 (2013); arXiv:1305.3342.
- [54] J. Kuipers, Q. Hummel and K. Richter, "Quantum graphs whose spectra mimic the zeros of the Riemann zeta function", *Phys. Rev. Lett.* **112**, 070406 (2014); arXiv:1307.6055.
- [55] G. Sierra, "The Riemann zeros as energy levels of a Dirac fermion in a potential built from the prime numbers in Rindler spacetime", *J. Phys. A: Math. Theor.* **47**, 325204 (2014); arXiv:1404.4252.
- [56] E. Yasar and Y. Yildirim, "A procedure on the first integrals of second-order nonlinear ordinary differential equations", *Eur. Phys. J. Plus* **130**, 240 (2015).
- [57] T. Okazaki, " AdS_2/CFT_1 , Whittaker vector and Wheeler-De Witt equation", *Phys. Rev. D* **92**, 126010 (2015); arXiv:1510.04759.
- [58] D. Schumayer, D. A. W. Hutchinson, "Physics of the Riemann Hypothesis", *Rev. Mod. Phys.* **83**, 307 (2011); arXiv:1101.3116.

- [59] E. Elizalde, V. Moretti, S. Zerbini, "On recent strategies proposed for proving the Riemann hypothesis", *Int. J. Mod. Phys. A* **18**, 2189 (2003); arXiv:math-ph/0109006.
- [60] H. C. Rosu, "Quantum hamiltonians and prime numbers", *Mod. Phys. Lett. A* **18**, 1205 (2003); arXiv:quant-ph/0304139.
- [61] M. Wolf, "Will a physicists prove the Riemann Hypothesis?", arXiv:1410.1214.
- [62] M. Watkins, *Number Theory and Physics* arxiv: <http://empslocal.ex.ac.uk/people/staff/mrwatkin//zeta/physics.htm>
- [63] M. du Sato, *The Music of the Primes*, Harper Collins, NewYork (2003).
- [64] K. Sabbagh, *Dr. Riemann's zeros*, Atlantic books, London (2003).
- [65] J. Derbyshire, *Prime obsession*, A plume book, New York (2003).
- [66] D. Rockmore, *Stalking the Riemann hypothesis*, Jonathan Cape, London (2005).
- [67] B. S. Pavlov and L. D- Fadeev, "Scattering theory and authomorphic functions", *Sov. Math.* **3**, 522 (1975). Plenum Publishing Corp. translation, N.Y;
- [68] P. D. Lax and R.S. Phillips, *Scattering Theory for Automorphic Functions*, Princeton University Press (1976).
- [69] R.K. Bhaduri, Avinash Khare, and J. Law, "Phase of the Riemann zeta function and the inverted harmonic oscillator", *Phys. Rev.* **E 52**, 486 (1995); chao-dyn/9406006.
- [70] A. Knauf, "Number theory, dynamical systems and statistical mechanics", *Reviews in Mathematical Physics.* **11**, 1027 (1999).
- [71] N. N. Khuri, Inverse scattering, the coupling constant spectrum, and the Riemann hypothesis, *Math. Phys. Anal. Geom.* **5**, 1 (2002); hep-th/0111067.
- [72] S. Joffily, Jost function, prime numbers and Riemann zeta function, math-ph/0303014.
- [73] A. LeClair, "Interacting Bose and Fermi gases in low dimensions and the Riemann hypothesis", *Int. J. Mod. Phys. A* **23**, 1371 (2008); arXiv:math-ph/0611043.
- [74] Y.-H. He, V. Jejjala, and D. Minic, Eigenvalue density, Lis positivity, and the critical strip, arXiv:0903.4321; Y.-H. He, V. Jejjala, and D. Minic, On the physics of the Riemann zeros, in the Proceedings of QTS6, Lexington, Kentucky, USA, July 20-25 2009, *J. Phys. Conf. Ser.* **462** (2013), S. R. Das and A. D. Shapere, eds.; arXiv:1004.1172.
- [75] E. de Rafael, "Large N_c QCD, Harmonic Sums and the Riemann Zeros", *Nucl. Phys. Proc. Suppl.* **207**, 290 (2010); arXiv:1010.4657.
- [76] S. L. Cacciatori and M. Cardella, Equidistribution rates, closed string amplitudes, and the Riemann hypothesis, *JHEP* **1012**, 025 (2010); arXiv:1007.3717;
- [77] C. Angelantonj, M. Cardella, S. Elitzur, and E. Rabinovici, Vacuum stability, string density of states and the Riemann zeta function, *JHEP* **1102**, 024 (2011); arXiv:1012.5091.
- [78] M.-A. Sanchis-Lozano, J. F. Barbero G., J. Navarro-Salas "Prime numbers, quantum field theory and the Goldbach conjecture", *Int. J. Mod. Phys. A* **27**, 1250136 (2012); arXiv:1201.6541.
- [79] M.V. Berry, "Riemann zeros in radiation patterns", *J. Phys. A: Math. Theor.* **45**, 302001 (2012).
- [80] M.V. Berry, "Riemann zeros in radiation patterns: II. Fourier transforms of zeta", *J. of Phys. A: Math. and Theor.* **48**, 385203 (2015).
- [81] J. I. Latorre and G. Sierra, "Quantum Computation of Prime Number Functions", *Quant. Inf. and Comp.*, **14**, 0577 (2014); arXiv:1302.6245.
- [82] G. Menezes, B. F. Svaiter and N. F. Svaiter, "Riemann zeta zeros and prime number spectra in quantum field theory", *Int. J. Mod. Phys. A* **28**, 1350128 (2013); arXiv:1303.7028.
- [83] R. V. Ramos and F. V. Mendes, "Riemannian Quantum Circuit", *Physics Letters A* **378** (2014) 1346 arXiv:1305.3759.
- [84] J. G. Dueñas and N. F. Svaiter, "Riemann zeta zeros and zero-point energy", *Int. J. Mod. Phys. A* **29**, (2014); arXiv:1311.6681.
- [85] C. Feiler and W.P. Schleich, "Entanglement and analytical continuation: an intimate relation told by the Riemann zeta function", *New J. Phys* **15**, 063009 (2013).
- [86] J. G. Dueñas, N. F. Svaiter, "Thermodynamics of the bosonic randomized Riemann gas". *J. Phys. A: Math. and Th.* **48**, 315201 (2015); arXiv:1401.8190.
- [87] J. I. Latorre and G. Sierra, "There is entanglement in the primes", *Quant. Inf. and Comp.*, **15**, 0622 (2015); arXiv:1403.4765.
- [88] J. G. Dueñas, N. F. Svaiter and G. Menezes, "One-loop effective action and the Riemann Zeros", *Int. J. of Mod. Phys. A* **29**, . (2014); arXiv:1405.6458.
- [89] C.E. Creffield, G. Sierra, "Finding zeros of the Riemann zeta function by periodic driving of cold atoms", *Phys. Rev. A* **91**, 063608 (2015); arXiv:1411.0459.
- [90] Yang-Hui He, Vishnu Jejjala, Djordje Minic, "From Veneziano to Riemann: A String Theory Statement of the Riemann Hypothesis", arXiv:1501.01975.
- [91] G. Franca and A. LeClair. "Transcendental equations satisfied by the individual zeros of Riemann, Dirichlet and modular L-functions". *Comm. in Number Theory and Physics*, **9**, No. 1 (2015); arXiv:1502.06003.
- [92] M. L. Lapidus, "The Sound of Fractal Strings and the Riemann Hypothesis", arXiv:1505.01548.
- [93] J. W. Neuberger, C. Feiler, H. Maier and W. P. Schleich, "The Riemann hypothesis illuminated by the Newton flow of ζ^* ", *Physica Scripta* **90**, 108015 (2015).
- [94] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions*, Dover, New York 1974. (1972).
- [95] W. G. Unruh, "Notes on black-hole evaporation", *Phys. Rev. D* **14**, 870 (1976).
- [96] J. von Neumann. *Math. Ann.*, **102**, 49 (1929).
- [97] A. Galindo and P. Pascual, "Quantum Mechanics I", Springer-Verlag, Berlin, 1991.
- [98] W. Rindler, "Kruskal space and the uniformly accelerated frame", *Am. J. Phys.* **34**, 1174 (1966).

- [99] G. Pólya, "Bemerkung über die integraldarstellung der Riemannschen zeta-funktion", *Acta Math.* **48**, 305 (1926).
 - [100] D. A. Hejhal, "On a result of G. Pólya concerning the Riemann ζ -function", *J. d Analyse Mathématique* **55**, 59 (1990).
 - [101] M. Asorey, A. Ibort, G. Marmo, "Global Theory of Quantum Boundary Conditions and Topology Change", *Int. J. Mod. Phys.* **A20**, 1001 (2005);hep-th/0403048.
 - [102] B. Julia, "Statistical theory of numbers, in Number Theory and Physics", eds. J. M. Luck, P. Moussa, and M. Waldschmidt, Springer Proceedings in Physics, Vol. 47, 276, Springer-Verlag, Berlin, 1990
 - [103] D. Spector, "Supersymmetry and the Moebius Inversion Function", *Commun. Math. Phys.* **127**, 239 (1990).
 - [104] G. Mussardo, "The quantum mechanical potential for the prime numbers"; cond-mat.9712010.
 - [105] H. Wu and D. W. L. Sprung, "Riemann zeros and a fractal potential", *Phys. Rev. E* **48**, 2595 (1993).
 - [106] A. Ramani, B. Grammaticos, and E. Caurier, "Fractal potentials from energy levels", *Phys. Rev. E* **51**, 6323 (1995).
 - [107] B. P. van Zyl and D. A. W. Hutchinson, "Riemann zeros, prime numbers, and fractal potentials", *Physical Review E*, **67**, 066211 (2003).
 - [108] D. Schumayer, B. P. van Zyl, P. Brandon and D. A. W. Hutchinson, "Quantum mechanical potentials related to the prime numbers and Riemann zeros", *Phys. Rev. E* **78**, 056215 (2008).
 - [109] J. Arias de Reyna and J. van de Lune, "On the exact location of the non-trivial zeros of Riemann's zeta function", arXiv:1305.3844.
 - [110] S. Blanes, F. Casas, J.A. Oteo and J. Ros "The Magnus expansion and some of its applications", *Phys. Rep.* **470**, 151 (2008).
 - [111] https://en.wikipedia.org/wiki/Mertens_conjecture
 - [112] A. M. Odlyzko, A. M. and H. J. J. te Riele, "Disproof of the Mertens conjecture", *Journal für die reine und angewandte Mathematik* **357**, 138 (1985).
 - [113] Nathan Ng, "The Distribution of the Summatory Function of the Möbius Function", *Proc. London Math. Soc.* **89**, 361 (2004).
 - [114] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer, New-York, 1976.
 - [115] R. Baillie, "Experiments with Zeta zeros and Perron's formula", arXiv:1103.6226.
 - [116] The expression for $\text{Re}_{s=0} F(s)$ corresponding to the case $m = 1$ contains the term $-\frac{1}{2}\zeta''(z)/(\zeta'(z))^2$ that was absent in reference [55]. However its addition does not modify the results obtained in [55].
-